arXiv:1211.4229v1 [math.CO] 18 Nov 2012

An upper bound on the fractional chromatic number of triangle-free subcubic graphs

Chun-Hung Liu*
School of Mathematics, Georgia Institute of Technology, Atlanta, Georgia 30332, USA

November 20, 2012

Abstract

An $(a:b)$-coloring of a graph G is a function f which maps the vertices of G into b-element subsets of some set of size a in such a way that $f(u)$ is disjoint from $f(v)$ for every two adjacent vertices u and v in G. The fractional chromatic number $\chi_f(G)$ is the infimum of a/b over all pairs of positive integers a, b such that G has an $(a:b)$-coloring. Heckman and Thomas conjectured that the fractional chromatic number of every triangle-free graph G of maximum degree at most three is at most 2.8. Hatami and Zhu proved that $\chi_f(G) \leq 3-3/64 \approx 2.953$. Lu and Peng improved the bound to $\chi_f(G) \leq 3-3/43 \approx 2.930$. Recently, Ferguson, Kaiser and Král' proved that $\chi_f(G) \leq 32/11 \approx 2.909$. In this paper, we prove that $\chi_f(G) \leq 43/15 \approx 2.867$.

1 Introduction

A *proper k-coloring* of a graph G is an assignment of one color of k colors to each vertex v of G such that adjacent vertices receive different colors. The *chromatic number* $\chi(G)$ of a graph G is the minimum k such that G has a proper k-coloring. Graph coloring is one of the most celebrated topics in graph theory, it has been widely explored and has many generalizations. For any positive integer a, we let $[a]$ be the set $\{1, 2, ..., a\}$. Given a set S, we define 2^S to be the collection of subsets of S. An *$(a:b)$-coloring* of a graph G is a function $f : V(G) \rightarrow 2^{[a]}$ such that $|f(v)| = b$ for every vertex v, and $f(x) \cap f(y) = \emptyset$ for every adjacent vertices x and y. The *fractional chromatic number* $\chi_f(G)$

---

*E-mail:cliu87@math.gatech.edu.

of G is the infimum of a/b over all pairs of positive integers a, b such that G has an $(a : b)$-coloring. In particular, every $(k : 1)$-coloring is a proper k-coloring, so $\chi_f(G) \leq \chi(G)$.

Fractional coloring can be investigated from the point of view of optimization. An *independent set* I of a graph is a subset of vertices such that every two vertices in I are not adjacent. So it is clear that the chromatic number is the minimum number of nonempty independent sets that partition the set of vertices. In other words, the chromatic number is the optimal value of a certain integer programming problem. In fact, the optimal value of the LP-relaxization of this integer programming problem is the fractional chromatic number, and the infimum in the definition of the fractional chromatic number is attained [13], so the infimum can be replaced by minimum. Furthermore, by taking advantage of the duality of linear programming problems, the fractional chromatic number of a graph G is at most k if and only if for every weighted function defined on vertices of G, there exists an independent set I such that the sum of weights of verties in I is at least w/k, where w is the sum of weights of all vertices in G.

Brooks' Theorem implies that $\chi_f(G) \leq \Delta(G)$, where $\Delta(G)$ is the maximum degree of the graph G unless G is the complete graph or an odd cycle. On the other hand, it is easy to show that $\chi_f(G) \geq \omega(G)$, where $\omega(G)$ is the *clique number* which is the maximum size of a subgraph whose vertices are pairwise adjacent in G, so one might expect a better upper bound when $\omega(G) < \Delta(G)$. Indeed, it is not hard to see that $\chi_f(G) \geq |V(G)|/\alpha(G)$, where $\alpha(G)$ is the maximum size of an independent set in G, so the best upper bound we can expect is $|V(G)|/\alpha(G)$.

A graph is *subcubic* if it has the maximum degree at most three. Given a family $\mathcal{H}$ of graphs, a graph is *$\mathcal{H}$-free* if it does not contain any graph in $\mathcal{H}$ as a subgraph. A graph is *triangle-free* if it is $\{K_3\}$-free. Staton [14] proved that $\alpha(G) \leq 5|V(G)|/14$ for every triangle-free subcubic graph G. This result is best possible as Fajtlowicz [3] pointed out that the generalized Petersen graph $P(7, 2)$ has 14 vertices but no independent set of size 6. Heckman and Thomas [7] gave a short proof of Staton's Theorem and gave the following conjecture.

Conjucture 1.1 *[7] The fractional chromatic number of every triangle-free subcubic graph is at most* $14/5$.

Some progress on this conjecture has been made, but it remains open. Let G be a triangle-free subcubic graph. Hatami and Zhu [6] proved that $\chi_f(G) \leq 3 - 3/64 \approx 2.953$, and $\chi_f(G) < 14/5$ if the shortest cycle of G has length at least 7. Lu and Peng [11] improved the bound to $\chi_f(G) \leq 3 - 3/43 \approx 2.930$. Recently, Ferguson, Kaiser and Král' [4] proved that $\chi_f(G) \leq 32/11 \approx 2.909$. In this paper, we prove the following theorem to improve the upper bound.

Theorem 1.2 *The fractional chromatic number of every triangle-free subcubic graph is at most* $43/15$.

Note that $43/15 = 14/5 + 1/15 \approx 2.867$.

The paper is organized as follows. We will give many lemmas without proofs and use them to prove Theorem 1.2 in section 2, and the remaining sections are dedicated to proving those lemmas. More specifically, we investigate the structure of fractionally critical graphs, which play the role of minimum counterexamples of Theorem 2 in section 3. In fact, we prove a more general result than we need in this paper, one that might be useful in future work. In section 4, we study a list-version of fractional colorings for graphs having some "simple" structures. In section 5, we prove the lemmas stated in section 2. Finally, we make some concluding remarks in section 6.

In the rest of this section, we introduce some terminology. In this paper, graphs do not contain multiple edges or loops. For any subset S of vertices of a graph $G = (V(G), E(G))$, we define the *neighborhood* $N_G(S)$ of S to be the set of vertices v which are not in S but are adjacent to a vertex in S. Also, we define $N_G[S]$ to be the set $N_G(S) \cup S$. And we write $N_G(\{v\})$ as $N_G(v)$ and $N_G[\{v\}]$ as $N_G[v]$ for short. If u, v are two vertices in G, then $G + uv$ is the graph obtained from G by adding an edge uv. If uv is an edge in G, then G/uv is the graph obtained from G by contracting uv and deleting resulting loops and parallel edges. We define $G[S]$ to be the subgraph induced by the set S when S is a subset of $V(G)$. The *degree* of a vertex v in G, denoted by $\deg_G(v)$, is the number of edges incident with v. G is *cubic* if every vertex in G is of degree 3. A *leaf* is a vertex of degree one, and a *support vertex* is a vertex that is adjacent to a leaf. A *matching* M is a subset of edges such that no two edges in M have a common end; we say that M *saturates* a vertex v if v is an end of some edge in M. Given any set S, we define the function 1_S by letting $1_S(x) = 1$ if $x \in S$, and $1_S(x) = 0$ for otherwise.

A *digraph* D is a graph equipped with orientation of edges. For each vertex v of D, we define $N_D^+(v)$ (and $N_D^-(v)$, repsectively) to be the set of vertices with head (tail, respectively) of an edge whose tail (head, respectively) is v. The *out-degree* $\deg_D^+(v)$ (*in-degree* $\deg_D^-(v)$, respectively) of v is the size of $|N_D^+(v)|$ ($|N_D^-(v)|$, respectively).

Let H be a subgraph of a graph G. We say a function f defined on $V(H)$ can be *extended* to a function g defined on $V(G)$ if $g(v) = f(v)$ for all $v \in V(H)$. We also say that f can be extended to G in this case.

2 Proof of Theorem 1.2

Given a graph G and a function $F : V(G) \to 2^{[14]}$, we say that $f : V(G) \to 2^{[14]}$ is an *F-avoiding coloring* if $f(v)$ is disjoint from $F(v) \cup f(u)$ for every two adjacent vertices u and v.

Let f be a proper 3-coloring of a subcubic graph G, $1 \leq i < j \leq 3$, and C a union of some components of the subgraph $G_{i,j}$ of G induced by $f^{(-1)}(i) \cup f^{(-1)}(j)$. For every vertex v in $N_G(C)$, we define $n_C(v)$ to be the number of vertices in C adjacent to v. We say that B is a *boundary-graph* of C if B is a graph such that $V(B) = N_G(C)$ and no three vertices x_1, x_2, x_3 with $n_C(x_i) = 1$ for $1 \leq i \leq 3$ form a triangle in B. $(B, \mathcal{J})$ is a *boundary-pair* of C if B is a boundary-graph of C and $\mathcal{J}$ is a collection of pairwise disjoint subsets J of $V(B)$ such that every vertex v in J satisfies that $n_C(v) \geq 2$. Given a boundary-pair $(B, \mathcal{J})$, we say that a function $F : V(C) \to 2^{[14]}$ is *$(B, \mathcal{J})$-compatible* if there is a $(14:2)$-coloring h of B such that each set $S \in \mathcal{J}$ contains two vertices x_S, y_S such that $h(x_S) = h(y_S)$, and $F(v) = \bigcup_{u \in N_G(v) \cap V(B)} h(u)$ for every vertex v of C. Note that $|F(v)| \leq 2|N_G(v) \cap V(B)| = 6 - 2\deg_C(v)$ for every vertex v in C and every $(B, \mathcal{J})$-compatible F.

We say that a boundary-pair $(B, \mathcal{J})$ of C *penetrates* C if there is an independent set $I \subseteq \{v \in V(C) : \deg_C(v) = 3\}$ in C such that for every $(B, \mathcal{J})$-compatible F, there are F-avoiding colorings g_1, g_2 of C such that $|g_1(v)| + |g_2(v)| = 16 - 2\deg_C(v) - 2 \cdot 1_I(v)$ for every vertex v of C. We say that a boundary-pair $(B_1, \mathcal{J}_1)$ of C *cooperates* with a boundary-pair $(B_2, \mathcal{J}_2)$ in C if there exists an independent set $I \subseteq \{v \in V(C) : \deg_C(v) = 3\}$ of C such that for every $k = 1, 2$ and $(B_k, \mathcal{J}_k)$-compatible F_k, there are F_k-avoiding colorings g_k of C such that $|g_1(v)| + |g_2(v)| = 16 - 2\deg_C(v) - 2 \cdot 1_I(v)$ for every vertex v of C.

Given two positive integers a, b, a set S, and two functions $f : S \to 2^{[a]}$ and $g : S \to 2^{[b]}$, define $f \uplus g : S \to 2^{[a+b]}$ by letting $f \uplus g(v) = \{x, y + a : x \in f(v), y \in g(v)\}$. Note that the operation $\uplus$ is not associative. For convenience, we simply denote $(...((f_1 \uplus f_2) \uplus f_3) \uplus ...) \uplus f_n$ by $f_1 \uplus f_2 \uplus f_3 \uplus ... f_n$ for every $n \geq 3$. The following lemma shows the motivation for introducing the notion of penetrations and cooperations.

Lemma 2.1 *Let f be a proper 3-coloring of G, $1 \leq s < t \leq 3$, p an integer, and let $C_1, C_2, ..., C_{r_{s,t}}$ be the components of the subgraph $G_{s,t}$ indeueced by $f^{(-1)}(s) \cup f^{(-1)}(t)$. Assume that there are boundry-pairs $(B_i, \mathcal{J}_i)$ that penetrates C_i for $1 \leq i \leq p$, and there are boundry-pairs $(B_{i,1}, \mathcal{J}_{i,1})$ and $(B_{i,2}, \mathcal{J}_{i,2})$ for $p+1 \leq i \leq r_{s,t}$ such that $(B_{i,1}, \mathcal{J}_{i,1})$ cooperates $(B_{i,2}, \mathcal{J}_{i,2})$ for $p+1 \leq i \leq r_{s,t}$. Let $R_k = \bigcup_{1 \leq i \leq p} B_i \cup$*

$\bigcup_{p+1\le i\le r_{s,t}} B_{i,k}$ *and* $\mathcal{S} = \bigcup_{1\le i\le p} \mathcal{J}_i \cup \bigcup_{p+1\le i\le r_{s,t}} \mathcal{J}_{i,k}$ *for* $k = 1, 2$. *If there are a* $(R_1, \mathcal{S}_1)$*-compatible function and a* $(R_2, \mathcal{S}_2)$*-compatible function, then there are an independent set* $I_{s,t} \subseteq \{v \in V(G_{s,t}) : \deg_{G_{s,t}}(v) = 3\}$ *and a function* $g_{s,t} : V(G) \to 2^{[56]}$ *such that* $|g_{s,t}(v)| = 8$ *if* $f(v) \notin \{s,t\}$, *and* $|g_{s,t}(v)| = 32 - 4\deg_{G_{s,t}}(v) - 4 \cdot 1_{I_{s,t}}(v)$ *if* $f(v) \in \{s,t\}$, *and* $g_{s,t}(x) \cap g_{s,t}(y) = \emptyset$ *for every adjacent vertices* x *and* y. *In particular, if such* $g_{1,2}, g_{1,3}$ *and* $g_{2,3}$ *exist, then* G *has a* $(172 : 60)$*-coloring, and* $\chi_f(G) \le 43/15$.

Proof. Let F_1 and F_2 be a $(R_1, \mathcal{S}_1)$-compatible and a $(R_2, \mathcal{S}_2)$-compatible function, respectively. Since $(B_i, \mathcal{J}_i)$ penetrates C_i for $1 \le i \le p$, there are independent sets $I_{s,t,i} \subseteq \{v \in V(C_i) : \deg_{G_{s,t}}(v) = 3\}$, F_1-avoiding colorings $h_{i,1}$ and $h'_{i,1}$, and F_2-avoiding colorings $h_{i,2}$ and $h'_{i,2}$ such that $|h_{i,1}(v)| + |h'_{i,1}(v)| = |h_{i,2}(v)| + |h'_{i,2}(v)| = 16 - 2\deg_{G_{s,t}}(v) - 2 \cdot 1_{I_{s,t,i}}(v)$ for vertex v in C_i and for $1 \le i \le p$. Similarly, since $(B_{i,1}, \mathcal{J}_{i,1})$ cooperates with $(B_{i,2}, \mathcal{J}_{i,2})$ for $p + 1 \le i \le r_{s,t}$. there are independent sets $I_{s,t,i} \subseteq \{v \in V(C_i) : \deg_{G_{s,t}}(v) = 3\}$ and F_k-avoiding colorings $h''_{i,k}$, for $k = 1, 2$, such that $|h''_{i,1}(v)| + |h''_{i,2}(v)| = 16 - 2\deg_{G_{s,t}}(v) - 2 \cdot 1_{I_{s,t,i}}(v)$ for vertex v in C_i and for $p+1 \le i \le r_{s,t}$.

Let $I_{s,t} = \bigcup_{1\le i\le r_{s,t}} I_{s,t,i}$, and let h_1, h_2 be a $(14 : 2)$-coloring of $(R_1, \mathcal{S}_1)$ and $(R_2, \mathcal{S}_2)$, respectively, such that for each $S \in \mathcal{S}_i$, there exist two vertices in S with the same h_i-value. Define $g_{s,t}(v) = (h_1 \uplus h_1 \uplus h_2 \uplus h_2)(v)$ for $v \in V(G) - V(G_{s,t})$, $g_{s,t}(v) = (h_{i,1} \uplus h'_{i,1} \uplus h_{i,2} \uplus h'_{i,2})(v)$ for $v \in V(C_i)$ and $1 \le i \le p$, and $g_{s,t}(v) = (h''_{i,1} \uplus h''_{i,1} \uplus h''_{i,2} \uplus h''_{i,2})(v)$ for $v \in V(C_i)$ and $p + 1 \le i \le r_{s,t}$. It is clear that $I_{s,t}$ is an independent set, and $g_{s,t}$ maps any two adjacent vertices to two disjoint sets in $2^{[56]}$. In addition, $|g_{s,t}(v)| = 8$ for $v \in V(G) - V(G_{s,t})$ and $|g_{s,t}(v)| = 32 - 4\deg_{G_{s,t}}(v) - 4 \cdot 1_{I_{s,t}}(v)$ for every vertex v in $G_{s,t}$.

If $g_{1,2}, g_{1,3}$ and $g_{2,3}$ exist, then define $g = g_{1,2} \uplus g_{1,3} \uplus g_{2,3}$. Hence, $g(v) \subseteq [168]$ and $|g(v)| = 72 - 4\deg_G(v) - 4(1_{I_{1,2}}(v) + 1_{I_{1,3}}(v) + 1_{I_{2,3}}(v))$ for every vertex v in G. Let $I = I_{1,2} \cup I_{1,3} \cup I_{2,3}$. Note that every vertex in $I_{1,2}$, $I_{1,3}$ or $I_{2,3}$ is adjacent to three vertices of the same f-value, so these three sets are pairwise disjoint, and I is an independent set as well. Define $g' : V(G) \to 2^{[172]}$ by assigning $g'(v) = g(v)$ if $v \notin I$, and $g'(v) = g(v) \cup \{169, 170, 171, 172\}$ if $v \in I$. Consequently, g' is an $(172 : 60)$-coloring since the maximum degree of G is at most three, and hence $\chi_f(G) \le 172/60 = 43/15$. ■

Lemma 2.2 *Let* G *be a triangle-free subcubic graph, and* f *a proper* 3*-coloring of* G. *Let* C *be a component of a subgraph of* G *induced by two color classes of* f. *Assume that* C *is a path of order* k, *where* k *is odd. If* $k \ne 3$, *then there is a boundary-graph* B *of* C *such that* $\deg_B(v) \le 2n_C(v)$ *for every vertex* v *in* B, *and* $(B, \emptyset)$ *penetrates* C. *If* $k = 3$, *then there are graphs* B_1 *and* B_2 *such that* $\deg_{B_i}(v) \le 2n_C(v)$

for every vertex v in B_i and for every $i = 1, 2$, and $(B_1, \emptyset)$ cooperates with $(B_2, \emptyset)$.

Let L_0 be the graph obtained from adding two vertices to a path of 4 vertices such that each of the new vertices is adjacent to the both ends of the path. We say that the tuple $(v_1, v_2, v_3, v_4, u_1, u_2)$ of six distinct vertices of G *forms a* L_0 in G if these six vertices induce a graph isomorhpic to L_0 and $u_1v_1v_2v_3v_4u_1$ and $u_2v_1v_2v_3v_4u_2$ are the two 5-cycles in L_0. Notice that two different L_0's in G are not necessary vertex-disjoint.

Given any proper 3-coloring f of G, we say that $(v_1, v_2, v_3, v_4, u_1, u_2)$ is a *rainbow* L_0 with respect to f (or f*-rainbow*) if $(v_1, v_2, v_3, v_4, u_1, u_2)$ forms a L_0, and $f(u_1) = f(u_2) = \pi(1)$, $f(v_1) = f(v_3) = \pi(2)$, $f(v_2) = f(v_4) = \pi(3)$ for some permutation π of $\{1, 2, 3\}$.

For every even numbers i, j, where $i, j \geq 4$, we denote the path of order i and j by P_i and P_j, respectively, and we define $H_{i,j}$ to be the graph that is obtained from two disjoint paths P_i and P_j by adding an edge incident with one support vertex in P_i and one support vertex in P_j. The family $\mathcal{H}$ consists graphs $H_{i,j}$ for all even numbers i, j with $i, j \geq 4$.

Lemma 2.3 *Let G be a triangle-free subcubic graph, and f a proper 3-coloring of G such that there are no rainbow L_0's with respect to f. Let C be a component of a subgraph of G induced by two color classes of f. If C is a path of even order or it is a graph in $\mathcal{H}$, then there are boundry-pairs $(B_1, \mathcal{J}_1)$ and $(B_2, \mathcal{J}_2)$ of C such that*

1. *Both $(B_1, \mathcal{J}_1)$ and $(B_2, \mathcal{J}_2)$ penetrate C.*
2. *For each $i = 1, 2$, every vertex v except at most one, denoted by w_i, in B_i has $\deg_{B_i}(v) \leq 2n_C(v)$ and $\deg_{B_i}(w_i) = 2n_C(w_i) + 1$ (if w_i exists).*
3. *If both w_1 and w_2 exist, then either $n_C(w_1) = n_C(w_2) = 3$ or $\deg_{B_{3-i}}(w_i) \leq 2n_C(w_i) - 1$ for each $i = 1, 2$.*
4. *For every i in which w_i exists, there is a vertex w'_i in the component of B_i containing w_i such that $\deg_{B_i}(w'_i) \leq 2n_C(w'_i) - 1$.*
5. *Each $\mathcal{J}_i$ contains at most one set.*
6. *If $\mathcal{J}_i = \{J_i\} \neq \emptyset$, then*
    - *(a) $|\bigcup_{v \in J_i, n_C(v)=2} N_{B_i}(v)| \leq 2$.*
    - *(b) If $|J_C| = 3$, then either $|\bigcup_{v \in J_i, n_C(v)=2} N_{B_i}(v)| \leq 1$, or there is a vertex v in $V(B_i)$ adjacent to exactly two vertices in J_i. For the latter case, $N_{B_i}(v) \subseteq J_i$ when $n_C(v) = 1$.*

(c) *Either* J_i *is an independent set of size* 2 *such that one vertex* x *in* J_i *has* $n_C(x) = 3$*, or* $J_i = \{x, y, z\}$ *is an independent set of size* 3*.*

(d) *If there exists* $x \in J_i$ *with* $n_C(x) = 3$*, then* $|\bigcup_{v \in J_i} N_{B_i}(v)| \leq 3$*. In this case, if* C *is a path of even order and some support vertex is of degree* 2 *in* G*, then* $|\bigcup_{v \in J_i} N_{B_i}(v)| \leq 3$*.*

We say that a graph G is *good* if G is bipartite subcubic and the following hold:

(G1) No vertex of degree three is adjacent to three vertices of degree three in G.

(G2) For every two adjacent vertices x, y of degree three in G, either at least one of x, y is adjacent to a leaf, or each of x, y is adjacent to a support vertex of degree two.

(G3) If x, y, z are three vertices of degree three such that xyz is a path in G, and each x and z is adjacent to a leaf, and the neighbor y' of y other than x and z has degree two, then the neighbor y'_1 of y' other than y is a leaf.

(G4) If x, y, z are three vertices of degree three such that xyz is a path in G, $\deg_G(z_1) \geq 2$, and $\deg_G(z_2) \geq 2$, where $N_G(z) = \{z_1, z_2, y\}$, then z_1 or z_2 is a support vertex of degree two.

We say that a 3-coloring f of G is *good* if it is proper, and G contains no f-rainbow L_0's, and every two color classes of f induce a good graph.

Lemma 2.4 *Let* G *be a triangle-free subcubic graph, and* f *a good* 3*-coloring of* G*. Let* C *be a component of a subgraph of* G *induced by two color classes of* f*. If* C *is not a path nor a graph in* $\mathcal{H}$*, then there are two boundary-graphs* B_1 *and* B_2 *of* C *such that* $\deg_{B_i}(v) \leq 2n_C(v)$ *for each* $i = 1, 2$*, and* $(B_1, \emptyset)$ *cooperates with* $(B_2, \emptyset)$*.*

G is *fractionally* t*-critical* if $\chi_f(G) > t$ but $\chi_f(H) \leq t$ for every proper subgraph H of G. Readers who are familar with the notion of coloring critical graphs might notice that the equality in the above definition is on the side of subgraphs instead of on the side of G as in the definition of coloring critical graphs. Even though our definition of fractionally critical graphs is not consistent with coloring critical graphs, we think that it is more reasonable to define critical graph in this way when we deal with fractional coloring.

Lemma 2.5 *If* G *is a fractionally* t*-critical triangle-free subcubic graph with* $t \geq 8/3$*, then* G *has a good* 3*-coloring.*

The following lemma is proved in [1] and probably elsewhere.

Lemma 2.6 *[1] If G is a graph, then there is an orientation of G such that the in-degree and the out-degree of every vertex v is at least $\lfloor \deg_G(v)/2 \rfloor$.*

Now we are ready to prove Theorem 1.2.

Proof of Theorem 1.2: Suppose to the contrary, and let G be a fractionally 43/15-critical triangle-free subcubic graph. By Lemma 2.5, there exists a good 3-coloring f of G. Let $1 \leq s < t \leq 3$ and let $G_{s,t}$ be the subgraph of G induced by $f^{(-1)}(s) \cup f^{(-1)}(t)$. For every component C of $G_{s,t}$ isomorphic to a path of even order or a graph in $\mathcal{H}$, there exist boundry-pairs $(B_{C,1}, \mathcal{J}_{C,1})$ and $(B_{C,2}, \mathcal{J}_{C,2})$ of C satisfiying statements 1, 2, ..., 6 in Lemma 2.3. So for each $i = 1, 2$, there is at most one vertex, denoted by $w_{C,i}$ if it exists, satisfies that $n_C(w_{C,i}) \leq 2$ and $\deg_{B_{C,i}}(w_{C,i}) = 2n_C(w_{C,i})+1$. We call a component C' of $G_{s,t}$ isomorphic to a path of even order or a graph in $\mathcal{H}$ *dangerous* if both $w_{C',1}$ and $w_{C',2}$ exist. Note that $w_{C,i}$ could be equal to $w_{C',i'}$ for some component $C' \neq C$ of $G_{s,t}$ and some i'. Construct a graph A, where $V(A) = \{w_{C',1}, w_{C',2} : C' \text{ is a dangerous component}\}$ and $E(A) = \{w_{C',1}w_{C',2} : C' \text{ is a dangerous component}\}$. By Lemma 2.6, there is an orientation of A such that every vertex v in A has in-degree and out-degree at most $\lceil \deg_A(v)/2 \rceil$. Since we can exchange the name of $B_{C,1}$ and $B_{C,2}$, we may assume that every directed edge of A in the orientation is from $w_{C,1}$ to $w_{C,2}$. Note that every $w_{C,i}$ is adjacent to at most three components of $G_{s,t}$, so the maximum degree of A is at most three, and $|\deg_A^+(v) - \deg_A^-(v)| \leq 1$ for every vertex v in A.

On the other hand, for every component C of $G_{s,t}$ not isomorphic to a path of even order or a graph in $\mathcal{H}$, either there is a boundry-pair $(B_C, \emptyset)$ penetrates C, or there are boundary-pairs $(B_{C,1}, \mathcal{J}_{C,1})$ and $(B_{C,2}, \mathcal{J}_{C,2})$ of C such that $(B_{C,1}, \mathcal{J}_{C,1})$ cooperates with $(B_{C,2}, \mathcal{J}_{C,2})$, and $\deg_{B_C}(v) \leq 2n_C(v)$ (or $\deg_{B_{C,i}}(v) \leq 2n_C(v)$ for $i = 1, 2$, respectively) for every vertex v in B_C (or $B_{C,1}$ and $B_{C,2}$, respectively), by Lemmas 2.2 and 2.4. For each $i = 1, 2$, construct a graph H_i by defining $V(H_i) = N_G(G_{s,t})$ and $E(H_i) = \bigcup E(B_C) \cup \bigcup E(B_{C,i})$, where the first union runs through all components C of $G_{s,t}$ which is a path of order not three, and the second union runs through all other components C of $G_{s,t}$. Note that for every vertex $v \in H_i$, $\deg_{H_i}(v) \leq 2\sum_C n_C(v) + |\{C : v = w_{C,i}\}| - |\{C : v = w_{C,3-i}\}| \leq 2\deg_G(v) + |\deg_A^+(v) - \deg_A^-(v)| \leq 7$. Similarly, we define $\mathcal{J}_i = \bigcup \mathcal{J}_{C,i}$ for each $i = 1, 2$, where the union runs through all components C of $G_{s,t}$ which is a path of order not 3.

Note that it is sufficient to show that for each $i = 1, 2$, there is a proper 7-coloring $h_i : V(H_i) \to [7]$ of H_i such that every set in $\mathcal{J}_i$ has two vertices of the same h_i-value. If such colorings h_1, h_2 exist, then it is clear that $F_i : V(H) \to 2^{[14]}$ defined by $F_i(v) = \{2h_i(v) - 1, 2h_i(v)\}$ for every vertex v in $V(H_i)$ is $(H_i, \mathcal{J}_i)$-compatible. Since it is true for every $1 \leq s < t \leq 3$, Lemma 2.1 implies that $\chi_f(G) \leq 43/15$, contradicting G is a fractionally 43/15-critical graph.

Now we shall show that there is a proper 7-coloring h_i of H_i such that every set in $\mathcal{J}_i$ contains two vertices having the same h_i-value. The proof for H_1 and H_2 are the same, so we write H instead of H_1, H_2 for convenience, and let $\mathcal{J}_C$, B_C and w_C be the corresponding $\mathcal{J}_{C,i}$, $B_{C,i}$ and $w_{C,i}$, respectively, for every component C of $G_{s,t}$.

First, we show that H is proper 7-colorable. By Brooks' Theorem, it is sufficient to show that no component of H is isomorphic to K_8, since the maximum degree of H is at most 7. Suppose to the contrary, and let X be a component of H isomorphic to K_8. The following two claims are clear, and Claim 3 is followed from statement 4 in Lemma 2.3.

Claim 1: If $\deg_X(v) = 7$ then v is adjacent to at least one dangerous component C of $G_{s,t}$ such that $v = w_C$.

Claim 2: If $\deg_X(v) = 7$ and $\deg_{B_C}(v) \leq 2n_C(v) - 1$ for some C, then $n_C(v) = 1$ and there are two other dangerous components C', C'' such that $v = w_{C'} = w_{C''}$.

Claim 3: For each dangerous component C, there is a vertex u_C in the component of B_C containing w_C with $\deg_{B_C}(u_C) \leq 2n_C(u_C) - 1$.

For any two vertices x, y in X, if $x = w_C$ and $y \in N_{B_C}(x)$ such that $\deg_{B_C}(y) \leq 2n_{B_C}(y) - 1$, then we direct the edge xy from x to y. Note that the direction of such edge is well-defined, since if xy is an edge in $B_{C'}$ and $B_{C''}$ for two different dangerous components C' and C'', then it is only counted once in the union when we define $E(H)$, so $\deg(x) \leq 6$, contradicting X is isomorphic to K_8. Consider the subgraph H' of H induced by edges that are oriented in X. Note that every vertex in H' has out-degree at least 1 (by Claims 1 and 3) and in-degree at most 1 (by Claim 2). Furthermore, if the vertex has in-degree 1, then it has out-degree 2 by Claim 2. So the sum of out-degree of vertices in H' is greater than the sum of in-degree of vertices in H', a contradiction. Hence, H is 7-colorable.

Denote $\mathcal{J}_C$ by $\{J_C\}$ for every nonempty $\mathcal{J}_C$. Note that for every component C, C' of $G_{s,t}$, we have that every vertex v in J_C satisfies that $n_C(v) \geq 2$, and no triangles in $B_{C'}$ contains three vertices whose $n_{C'}$-values are 1, so J_C is not a clique in H. In addition, every vertex v in J_C has degree at most 6 in H.

Claim 4: Every proper 7-coloring of $H - J_C$ can be extended to a proper 7-coloring of H such that some two vertices in J_C receive the same color if one of the following is satisfied:

1. $|J_C| = 2$;
2. there are two edges in H with both ends in J_C;
3. there is a vertex in J_C incident with at most two edges in $E(H) - E(B_C)$, and there are at most one edge in H with both ends in J_C;
4. $|\bigcup_{u \in J_C} N_{B_c}(u)| = 1$.

Proof of Claim 4:

1. If $|J_C| = 2$, then let $J_C = \{x, y\}$. By statement 6 (c) and (d) in Lemma 2.3, we know that one vertex, say x, in J_C satisfies that $n_C(x) = 3$, $|N_{B_C}(x) \cup N_{B_C}(y)| \leq 3$ and $\deg_{H-B_C}(y) \leq 3$, so $|N_H(x) \cup N_H(y) - J_C| \leq 6$. Therefore, we can extend any proper 7-coloring of $H - J_C$ to H such that x and y receive the same color.
2. We may assume that $|J_C| = 3$ for the rest of the proof of Claim 4. Let $J_C = \{x, y, z\}$ and $xy, yz \in E(H)$. Since $n_C(v) \geq 2$ for every $v \in J_C$, there is at most one component $C' \neq C$ of $G_{s,t}$ such that $J_C \cap V(B_{C'}) \neq \emptyset$. Also, J_C is an independ set in B_C and x, y, z cannot form a triangle in $B_{C'}$, so $xz \notin E(H)$. In addition, $|N_H(x) \cup N_H(z) - J_C| \leq 6$, since $|N_{B_C}(x) \cup N_{B_C}(z)| \leq 2$ and both $\deg_{B_{C'}-\{xy\}}(x)$ and $\deg_{B_{C'}-\{xy\}}(y)$ are at most 2 for the component $C' \neq C$ of $G_{s,t}$ containing x, y (if C' exists), so every proper 7-coloring h of $H - J_C$ can be extended to H such that $h(x) = h(z)$ by first defining $h(x) = h(z)$ to be a color in $[7] - (N_H(x) \cup N_H(z) - J_C)$ and then defining $h(y)$ to be any feasible color.
3. Assume that there is a vertex in J_C incident with at most two edges in $E(H) - E(B_C)$. If there is exactly one edge in H with both ends in J_C, then there are two non-adjacent vertices u, v in J_C such that $|N_H(u) \cup N_H(v) - J_C| \leq 6$, so every proper 7-coloring of H can be extended to a proper 7-coloring of H such that two vertices in J_C receive the same color.

   So we may assume that no edge in H with both ends in J_C. Let $J_C = \{a, b, c\}$, and let a be incident with at most two edges in $E(H) - E(B_C)$. Then for every proper 7-coloring h of $H - J_C$, $|(N_H(a) \cup N_H(b)) - J_C|$ and $|(N_H(a) \cup N_H(c)) - J_C|$ are at most 7, so either $|h((N_H(a) \cup N_H(b)) - J_C)| \leq 6$ or $|h((N_H(a) \cup N_H(c)) - J_C)| \leq 6$ or $h(N_H(b) - J_C) = h(N_H(c) - J_C)$, so there are two non-adjacent vertices u, v in J_C such that $|h(N_H(u) \cup N_H(v) -$

$J_C)| \leq 6$, and hence it can be extended to a proper 7-coloring of H such that $h(u) = h(v)$.

4. If $|\bigcup_{u \in J_C} N_{B_c}(u)| = 1$, then we can use the similar argument in the last paragraph to show that for every proper 7-coloring of $H - J_C$, there are two vertices x, y in J_C such that $|h(N_H(x) \cup N_H(y) - J_C)| \leq 6$, so h can be extended to a proper 7-coloring of H such that $h(x) = h(y)$. This completes the proof of Claim 4. $\square$

For each component C of $G_{s,t}$, if any hypothesis of Claim 4 applies on J_C, then define $S_C = J_C$; otherwise, define $S_C = J_C \cup \{u_C\}$, where u_C is a vertex in $\bigcup_{u \in J_C} N_{B_C}(u)$ such that u_C is adjacent to exactly two vertices in J_C. Note that u_C exists by statement 6 in Lemma 2.3. By Claim 4, if $S_C = J_C$ and h is a proper 7-coloring of $H - J_C$, then h can be extended to a proper 7-coloring of H such that at least two vertices in J_C receive the same color. On the other hand, if $u_C \in S_C \cap J_{C'}$ for some two distinct components C and C', then $n_C(u_C) = 1$, $n_{C'}(u_C) = 2$, and there are at most two edges in $E(H) - E(J_{C'})$ incident with u_C, so $S_{C'} = J_{C'}$. Therefore, it is sufficient to find a proper 7-coloring of $H - \bigcup_C J_C$, where the union runs through all components C of $G_{s,t}$ such that $S_C = J_C$, such that $J_{C'}$ has two vertices receiving the same color for each component C' of $G_{s,t}$ with the property that $S_{C'} \neq J_{C'}$. Hence, we may assume that $S_C \neq J_C$ for every component C of $G_{s,t}$. Since every vertex v in J_C has $n_C(v) \geq 2$, we have that $S_C \cap S_{C'} \subseteq \{u_C, u_{C'}\}$ for every two different components C, C' of $G_{s,t}$. Using the argument for proving the fourth case of Claim 4, one can show that every 7-coloring h of $H - S_C$ can be extended to a proper 7-coloring of $(H - S_C) \cup \{x, y\}$ such that $h(x) = h(y)$ for some distinct $x, y \in J_C$. Furthermore, we can further extend the coloring h of $(H - S_C) \cup \{x, y\}$ to $(H - S_C) \cup \{x, y, u_C\}$, since u_C adjacent to at least two vertices in J_C, it means that either two neighbors of u_C have the same h-value, or one neighbor of u_C has not been assigned h-value. Finally, we can extend h to H since every vertex in J_C has degree at most 6.

Now, we constuct a proper 7-coloring h of H such that each J_C has two vertices getting the same h-value. Note that $J_C \cap J_{C'}$ unless $C = C'$. Set $H' = H - \bigcup_C S_C$, where the union runs through all components C of $G_{s,t}$ such that $\mathcal{J}_C \neq \emptyset$, and let h be a proper 7-coloring of $H - \bigcup_C S_C$. Pick a component C^*, where $\mathcal{J}_{C^*} = \{J_{C^*}\} \neq \emptyset$ and $S_{C^*} \neq J_{C^*}$ and $J_{C^*} \cap H' = \emptyset$, then extend h to $H' \cup J_{C^*}$ such that $h(x_{C^*}) = h(y_{C^*})$ for some distinct vertices x_{C^*} and y_{C^*} in J_{C^*}, and set H' to be the subgraph of H induced by $V(H') \cup J_{C^*}$. Repeat this process until no such C^* exists. Note that $J_{C^*} \cap S_C = \emptyset$ for other component C of $G_{s,t}$, since $J_{C^*} \neq S_{C^*}$. So any vertex in H that has not been colored by h is either in J_C with $J_C = S_C$ or equal to u_C with $J_C \neq S_C$. According to the augument in the previous paragraph, we can extend h to the remaining vertices in H. This completes the proof of the theorem. $\square$

3 Structure of fractionally critical graphs

The objective of this section is to prove Lemma 2.5. Recall that G is *fractionally t-critical* if $\chi_f(G) > t$ but $\chi_f(H) \leq t$ for every proper subgraph H of G. The following lemma is obvious.

Lemma 3.1 *Let $t > 0$. Every fractionally t-critical graph is 2-connected, and it has no vertex-cut which induces a clique.*

Lu and Peng [11] gave the following result for graphs that have a vertex-cut with size 2.

Lemma 3.2 *[11] Let G_1 and G_2 be two induced subgraph of G such that $V(G_1) \cup V(G_2) = V(G)$ and $V(G_1) \cap V(G_2) = \{u, v\}$, where $\{u, v\}$ is a vertex-cut.*

1. *If uv is an edge in G, then $\chi_f(G) = \max\{\chi_f(G_1), \chi_f(G_2)\}$.*
2. *If uv is not an edge in G, then $\chi_f(G) \leq \max\{\chi_f(G_1), \chi_f(G_2 + uv), \chi_f(G_2/uv)\}$.*

Now, we shall give a list of forbidden subgraphs for every fractionally t-critical graph for $t \geq 8/3$. We define R_i to be the graphs in Figure 1 for $0 \leq i \leq 7$.

Lemma 3.3 *Every fractionally t-critical graph with $t \geq 8/3$ is $\{R_i : 1 \leq i \leq 7\}$-free.*

Proof. Suppose that H is an induced subgraph of a fractionally t-critical graph G with $t \geq 8/3$ isomorphic to R_i for some $1 \leq i \leq 7$. It is clear that R_i has a $(8 : 3)$-coloring for every $1 \leq i \leq 7$, so $|N(H)| \geq 2$ by Lemma 3.1. Since R_5 and R_6 are cubic, and R_3 contains only one vertex of degree at most two, so $i \neq 3, 5, 6$. Let u_i and v_i be the two vertices in $N(H)$ when $i = 1, 2, 4, 7$, and let H' be the subgraph of G induced by $V(H) \cup \{u_i, v_i\}$. Note that u_i might be equal to v_i, and either $\{u_i, v_i\}$ is a vertex-cut or $H' = G$. It is easy to check that $\chi_f(H' + u_iv_i) \leq 8/3$ (when $u_i \neq v_i$) and $\chi_f(H'/u_iv_i) \leq 8/3$ for every $i = 1, 2, 4, 7$. Hence, every fractionally t-critical graph is R_i-free, for $i = 1, 2, 4, 7$, by Lemma 3.2 $\blacksquare$



Figure 1: Graphs R_i for $0 \leq i \leq 7$

Next, we investigate structure of fractionally t-critical graphs that contains L_0 as a subgraph, where $t \geq 8/3$. Note that two different L_0's are not necessarily vertex-disjoint. For $i \in \mathbb{N}$, define $\mathcal{L}_0 = \{L_0\}$ and $\mathcal{L}_i$ to be the collection of triangle-free subcubic graphs H that can be obtained from some graph H' in $\mathcal{L}_{i-1}$ by either

- (**Operation 1**) adding a disjoint P_2 to H' and two edges, where one edge is incident with one end of the P_2 and one vertex in a C_4 in H', and the other edge is incident with the other end of the P_2 and the diagonal vertex of the same C_4 in H', or
- (**Operation 2**) adding a disjoint C_4 to H' and two edges, where one edge is incident with one vertex in the C_4 and one end of a P_2 in H', and the other edge is incident with the diagonal vertex in the C_4 and the other end of the P_2.

It is easy to see that every graph in $\mathcal{L}_i$, where $i \geq 0$, contains exactly 4 vertices of degree 2, and these 4 vertices can be paired such that each pair either induces a path of order 2 or contains two diagonal vertices of a 4-cycle. For every finite sequence $(a_1, a_2, ..., a_k)$ with $a_i \in \{1, 2\}$ for all $1 \leq i \leq k$, we denote $L_{a_1,a_2,...,a_k}$ as the graph that is obtained from L_0 by doing Operations $a_1, a_2, ..., a_k$ consecutively. Note that for every graph $H \in \mathcal{L}_t$ and for each Operation j, where t is a postitive integer and $j = 1, 2$, there is at most one way (up to isomorphism) to add edges to link H and the new P_2 or C_4, so $L_{a_1,a_2,...,a_k}$ is well-defined. However, $L_{a_1,a_2,...,a_k}$ may be equal to $L_{b_1,b_2,...,b_m}$ for some two differnent sequences $(a_1, a_2, ..., a_k)$ and $(b_1, b_2, ..., b_m)$. The following lemma ensures that every graph in $\mathcal{L}_t$ can be generated by Operations 1 and 2 alternately, and it can be easily proved by doing induction on t, so we leave the proof for the readers.

Lemma 3.4 *Every graph H in $\mathcal{L}_t$, there is a sequence $(a_1, a_2, ..., a_t)$ with $a_1 \in \{1, 2\}$ and $a_{i+1} = 3 - a_i$ for $1 \leq i \leq t - 1$ such that $H = L_{a_1,a_2,...,a_t}$.*

We denote $L_{a_1,a_2,...,a_t}$ by $L^{(t)}_{a_1}$ for each sequence $(a_1, a_2, ..., a_t)$ with $a_i \in \{1, 2\}$ and $a_{i+1} = 3 - a_i$. By Lemma 3.4, $\mathcal{L}_t = \{L^{(t)}_1, L^{(t)}_2\}$ for each positive integer t. Let $\mathcal{L}'$ be the set of triangle-free subcubic graphs H' that contains a graph H in $\mathcal{L}_i$, for some i, as a spanning subgraph with $|E(H')| > |E(H)|$.

Lemma 3.5 *Every fractionally t-critical triangle-free subcubic graph with $t \geq 8/3$ is $\mathcal{L}'$-free.*

Proof. Let G be a fractionally t-critical triangle-free subcubic graph with $t \geq 8/3$. Since every graph in $\mathcal{L}_i$, where $i \geq 0$, has at most 4



Figure 2: A $(8:3)$-coloring of a subgraph of a graph in $\mathcal{L}_k$

vertices of degree two, every graph in $\mathcal{L}'$ has at most two vertices of degree two. Hence, if G is not $\mathcal{L}'$-free, then either G is in $\mathcal{L}'$, or there are two vertices u, v such that $G-\{u,v\}$ is not connected and one component of $G-\{u,v\}$ is in $\mathcal{L}'$, since G is 2-connected. We shall show that $\chi_f(G) \leq 8/3$, which leads to a contradiction.

For the case that G is in $\mathcal{L}'$, we assume that G contains a graph H in $\mathcal{L}_k$ as a spanning subgraph for some positive integer k, and we may assume that G is cubic. For the case that $G-\{u,v\}$ contains a subgraph in $\mathcal{L}'$, we let H be a component of $G-\{u,v\}$ containing a graph in $\mathcal{L}_k$, and it is sufficient to show that $\chi_f(H+uv)$ and $\chi_f(H/uv)$ are at most $8/3$ by Lemma 3.2. Let $\{w,x,y,z\}$ be the four vertices of degree two in H. So, we may assume one of the following cases holds by symmetry:

1. Both wx and yz are edges in H.
2. wx is an edge in H and y,z are diagonal vertices in a C_4.
3. w,x are diagonal vertices in a C_4 and y,z are diagonal vertices in another C_4.

Note that the neighbor of u and the neighbor of v are not adjacent in H since G is triangle-free and $|E(H)| > |E(H')|$ for $H' \in \mathcal{L}_k$

Observe that if f is a $(8:3)$-coloring of a graph $A-\{b,c\}$, where $P=abcd$ is a path in A such that internal vertices of P are of degree two in A, and $f(a) \neq f(d)$, then there is a $(8:3)$-coloring of A obtained from f by extending it to $V(A)$. On the other hand, if f is a $(8:3)$-coloring of a graph $A-\{b\}$, where $P=abc$ is a path in A such that the internal vertex b of P is of degree two in A, and $f(a)\cap f(c) \neq \emptyset$, then there is a $(8:3)$-coloring of A obtained from f by extending it to $V(A)$. Consequently, it is sufficient to show that:

1. If wx and yz are edges, then there are $(8:3)$-colorings f_1, f_2 of H such that $f_1(w)\cap f_1(y)=\emptyset$, $f_1(x)\cap f_1(z)=\emptyset$, $f_2(w)\cap f_2(y)=\emptyset$, and $1 \leq |f_2(x)\cap f_2(z)| \leq 2$.
2. If wx is an edge and y,z lie in a C_4, then there are $(8:3)$-colorings f_1, f_2 of H such that $f_1(w)\cap f_1(y)=\emptyset$, $f_1(x)\cap f_1(z)=$

$\emptyset$, $f_2(w) \cap f_2(y) = \emptyset$, $1 \leq |f_2(x) \cap f_2(z)| \leq 2$.

3. If w, x are diagonal vertices in a C_4 and y, z are diagonal vertices in a C_4, then there are $(8:3)$-colorings f_1, f_2 of H such that $f_1(w) \cap f_1(y) = \emptyset$, $f_1(x) \cap f_1(z) = \emptyset$, $f_2(w) \cap f_1(y) = \emptyset$, $1 \leq |f_2(w) \cap f_2(y)| \leq 2$.

The coloring drawn in Figure 2 shows that the patterns of the colors on the two adjacent degree-two vertices can be kept if we do Operations 2,1,2,1 consecutively, so it is sufficient to check the above three cases for $0 \leq k \leq 4$. In fact, it is not hard to check these cases, and we leave this for the readers. ■

Lemma 3.6 *Let G be a $\{K_3, R_i : 1 \leq i \leq 7\} \cup \mathcal{L}'$-free subcubic graph. Let t be a postitive integer, and for each i with $1 \leq i \leq t$, let H_i be a subgraph of G such that H_i is isomorphic to L_0. If $V(H_{i+1}) \cap (\bigcup_{j=1}^{i} V(H_j)) \neq \emptyset$ for all $i \geq 1$, then $G[\bigcup_{i=1}^{t} V(H_i)]$ is isomorphic to R_0 or a graph in $\mathcal{L}_k$ for some $k \geq 0$.*

Proof. For each $1 \leq i \leq t$, let $V(H_i) = \{v_{i,1}, v_{i,2}, v_{i,3}, v_{i,4}, u_{j,1}, u_{j,2}\}$, and $(v_{i,1}, v_{i,2}, v_{i,3}, v_{i,4}, u_{i,1}, u_{i,2})$ form L_0. We shall prove this lemma by doing induction on t. The lemma is clear if $t = 1$. Now, we prove the case that $t = 2$.

Suppose this lemma is not true when $t = 2$. If $(v_{2,1}, v_{2,2}) = (u_{1,1}, v_{1,1})$, then $G[V(H_1) \cup V(H_2)]$ is isomorphic to R_0 by the triangle-freeness. So we may assume that $(v_{2,1}, v_{2,2}) \neq (u_{1,1}, v_{1,1})$. If $v_{2,1} = u_{1,1}$, then $v_{2,2} \notin V(H_1)$ by symmetry, and hence $G[V(H_1) \cup V(H_2)]$ is isomorphic to the graph that can be obtained from L_0 by Operation 1. So we may assume that $\{v_{2,1}, v_{2,4}\} \cap \{u_{1,1}, u_{1,2}\} = \emptyset$. If $v_{2,2} = u_{1,1}$ and $v_{2,1} \in V(H_1)$, then $G[V(H_1) \cup V(H_2)]$ is isomorphic to the graph that can be obtained from L_0 by Operation 1 and then adding an edge, so it is in $\mathcal{L}'$. If $v_{2,2} = u_{1,1}$ and $v_{2,1} \notin V(H_1)$, then $G[V(H_1) \cup V(H_2)]$ is isomorphic to R_1, a contradition. Hence, we may assume that $\{u_{1,1}, u_{1,2}\} \cap \{v_{2,j} : 1 \leq j \leq 4\} = \emptyset$. Consequently, if $v_{2,1}$ or $v_{2,4}$ is in $V(H_1)$, say $v_{2,1} \in V(H_1)$, then $v_{2,1}$ is equal to $v_{1,1}$ or $v_{1,4}$, and hence $G[V(H_1) \cup V(H_2)]$ is isomorphic to L_0, R_3 or R_4. So $v_{2,1}$ and $v_{2,4}$ are not in $V(H_1)$. Furthermore, if $v_{2,2}$ or $v_{2,3}$ is in $V(H_1)$, say $v_{2,2} \in V(H_1)$, then $v_{2,2}$ is $v_{1,2}$ or $v_{1,3}$, and hence $G[V(H_1) \cup V(H_2)]$ is isomorphic the graph that can be obtained from L_0 by Operation 2. As a result, $\{v_{2,j} : 1 \leq j \leq 4\}$ is disjoint from $V(H_1)$, and it implies that $\{u_{2,1}, u_{2,2}\} \cap V(H_1) = \emptyset$, so $V(H_1) \cap V(H_2) = \emptyset$, a contradiction. This proves the case that $t = 2$.

Let $t \geq 3$. First, suppose that $G[\cup_{i=1}^{t-1} V(H_i)] = R_0$. If $\{v_{t,1}, v_{t,4}\} \cap (\bigcup_{i=1}^{t-1} V(H_i)) \neq \emptyset$, then $G[\cup_{i=1}^{t} V(H_i)]$ is isomorphic to R_0, R_1, R_2, R_3, R_5, R_6 or R_7, a contradiction. So $\{v_{t,1}, v_{t,4}\} \cap (\bigcup_{i=1}^{t-1} V(H_i)) = \emptyset$, but it implies that $\{v_{t,2}, v_{t,3}, u_{t,1}, u_{t,2}\} \cap (\bigcup_{i=1}^{t-1} V(H_i)) = \emptyset$ as well, a contradiction. Therefore, $G[\cup_{i=1}^{t-1} H_i]$ is isomorphic to a graph in $\mathcal{L}_k$ for some $k \geq 0$ by the induction hypothesis. We may assume that $k \geq 2$ since we may assume that $V(H_{i+1}) - (\bigcup_{j=1}^{i-1} V(H_j)) \neq \emptyset$ for $1 \leq i \leq t-1$, without loss of generality. In this case, we have the following two claims since G is $\mathcal{L}'$-free.

Claim 1: Each vertex of degree two in a graph in $\mathcal{L}_k$ is adjacent to a vertex of degree three.

Claim 2: Each two non-adjacent vertices of degree two in a graph in $\mathcal{L}_k$ either are in a 4-cycle or have distance at least three.

Observe that $\{v_{t,j} : 1 \leq j \leq 4\} \cap \bigcup_{j=1}^{t-1} V(H_j) \neq \emptyset$, since every graph in $\mathcal{L}_k$ has minimum degree two. So it is impossible that $v_{t,1}, v_{t,2} \notin V(H_t) - (\bigcup_{j=1}^{t-1} V(H_j))$ by Claims 1 and 2. Assume that $v_{t,1} \notin V(H_t) - (\bigcup_{j=1}^{t-1} V(H_j))$, but $v_{t,2} \in V(H_t) - (\bigcup_{j=1}^{t-1} V(H_j))$. By Claim 2, $v_{t,4} \notin V(H_t) - (\bigcup_{j=1}^{t-1} V(H_j))$, but it implies that $G[\bigcup_{j=1}^{t} V(H_j)] \in \mathcal{L}_{k+1}$. Hence, we may assume that $v_{t,1}, v_{t,4} \in \bigcup_{j=1}^{t-1} V(H_j)$ by symmetry. Therefore, one of $u_{t,1}$ and $u_{t,2}$ is in $\bigcup_{j=1}^{t-1} V(H_j)$ as well, so $v_{t,1}$ and $v_{t,4}$ have distance two in $\bigcup_{j=1}^{t-1} V(H_j)$, and this implies that one of $v_{t,1}$ and $v_{t,4}$ has degree three in $\bigcup_{j=1}^{t-1} V(H_j)$ by Claim 2, and hence both $u_{t,1}$ and $u_{t,2}$ are in $\bigcup_{j=1}^{t-1} V(H_j)$, and $v_{t,1}$ and $v_{t,2}$ are diagonal vertices in a 4-cycle. Consequently, $G[\bigcup_{j=1}^{t} V(H_j)]$ is isomorphic to a graph in $\mathcal{L}_{k+1}$. This completes the proof. ■

Let G be a graph and H a subgraph, then we say that a proper 3-coloring f of G is *H-rainbow-free* if H contains no rainbow L_0 with respect to f.

Lemma 3.7 *Let G be a subcubic $\{K_3\} \cup \mathcal{L}'$-free graph and H a subgraph of G isomorphic to a graph in $\mathcal{L}_t$ for some $t \geq 0$. Let f be a proper 3-coloring of $G - H$. If either $\deg_G(v) < 3$ for some vertex v in H, or $|f(N_G(H))| \geq 2$, then f can be extended to a proper H-rainbow-free 3-coloring of G.*

Proof. We shall prove this lemma by doing induction on t. We may assume that $\deg_G(x) = 3$ for all $x \in V(H)$, otherwise, for each vertex v in H of degree at most 2, we add a new vertex u_v to G and a new edge $u_v v$, and then define $f(u_v)$ to be an element in $\{1, 2, 3\}$ such that $|f(N_{G'}(H))|$ is as large as possible, where G' is the new graph, then $|f(N_{G'}(H))| \geq 2$, and any proper H-rainbow-free 3-coloring g of G' extended from f is a proper H-rainbow-free 3-coloring of G as well. So it is sufficient to deal with the case that $|f(N_G(H))| \geq 2$.

When $t = 0$, H is an induced subgraph as G is triangle-free. Let $V(H) = \{u_i, v_j : i = 1, 2, j = 1, 2, 3, 4\}$ and $(v_1, v_2, v_3, v_4, u_1, u_2)$ form a L_0. Let $x_i \in N_G(u_i) - V(H)$, and $y_j \in N_G(v_j) - V(H)$, for $i = 1, 2$ and $j = 2, 3$. Then f can be extended to a proper H-rainbow-free 3-coloring g of G by first defining $g(v_1) = g(v_4) = f(y_2)$ and then defining g on the remaining vertices in H.

Now, we assume that $t \geq 1$, so H is isomorphic to $L_k^{(t)}$ for some $k = 1, 2$. First, we assume that $k = 2$. So there is a cycle $abcda$ such that $H - \{a, b, c, d\}$ is isomorphic to a graph in $\mathcal{L}_{t-1}$, where $\deg_H(a) = \deg_H(c) = 2$. Let a', c' be the neighbor other than in $\{b, d\}$ of a and c, respectively. Let $\{x, y\} = N_G(H) - \{a', c'\}$. Note that a', c', x, y may not be pairwise distinct. If $f(a') \neq f(c')$, say $f(a') = 1$ and $f(c') = 2$, then without loss of generality, we may assume that $(f(x), f(y)) \neq (1, 1)$ (since we can swap colors 1 and 2), and hence we can extend f to a proper H-rainbow-free 3-coloring g of G by first defining $(g(a), g(b), g(c), g(d)) = (3, 1, 3, 1)$, and then defining g on $V(H) - \{a, b, c, d\}$ by the induction hypothesis. If $f(a') = f(c')$, say $f(a') = f(c') = 1$, then without loss of generality, we may assume that $(f(x), f(y)) \neq (3, 3)$ (since we can swap colors 2 and 3), and hence we can extend f to a proper H-rainbow-free 3-coloring g of G by first defining $(g(a), g(b), g(c), g(d)) = (2, 3, 2, 3)$, and then defining g on $V(H) - \{a, b, c, d\}$ by the induction hypothesis. This proves the case that $k = 2$.

Assume that $k = 1$. So there is a tuple $(p_1, p_2, p_3, p_4, q_1, q_2)$ of vertices in H such that $(p_1, p_2, p_3, p_4, q_1, q_2)$ forms a L_0 with $\deg_H(p_2) = \deg_H(p_3) = 2$, and $H - \{p_1, p_2, p_3, p_4, q_1, q_2\}$ is in $\mathcal{L}_{t-2}$ if $t \geq 2$, and a P_2 if $t = 1$. Let p'_2, p'_3 be the neighbor other than in $\{p_1, p_2, p_3, p_4\}$ of p_2, p_3, respectively. Let $\{x, y\} = N_G(H) - \{p_2, p_3\}$. Note that p'_2, p'_3, x, y may not be pairwise distinct. Suppose that $t \geq 2$. If $f(p'_2) \neq f(p'_3)$, say $f(p'_2) = 1, f(p'_3) = 2$, then without loss of generality, we may assume $(f(x), f(y)) \neq (1, 1)$ (since we can swap colors 1 and 2), and hence we can extend f to a proper H-rainbow-free 3-coloring g of G by first defining $(g(p_1), g(p_2), g(p_3), g(p_4), g(q_1), g(q_2)) = (2, 3, 1, 3, 1, 1)$ and then defining g on $V(H) - \{p_1, p_2, p_3, p_4, q_1, q_2\}$ by the induction hypothesis. If $f(p'_2) = f(p'_3)$, say $f(p'_2) = f(p'_3) = 1$, then without loss of generality, we may assume $(f(x), f(y)) \neq (2, 2)$ (since we can swap colors 2 and 3), and hence we can extend f to a proper H-rainbow-free 3-coloring g of G by first defining $(g(p_1), g(p_2), g(p_3), g(p_4), g(q_1), g(q_2)) = (3, 2, 3, 1, 2, 2)$ and then defining g on $V(H) - \{p_1, p_2, p_3, p_4, q_1, q_2\}$ by the induction hypothesis. This proves the case that $k = 2$ and $t \geq 2$.

As for the case that $k = 2$ and $t = 1$, note that the above arguments also work except for the case that $f(x) = f(y)$. By symmetry, it is sufficient to consider the case that $f(p'_2) = f(p'_3)$ and $f(x) = f(y)$. Without loss of generality, we may assume that $f(p'_2) = f(p'_3) = 1$ and

$f(x)=f(y)=2$. Then it is clear that we can extend f to a proper H-rainbow-free 3-coloring g by defining $(g(p_1),g(p_2),g(p_3),g(p_4),g(q_1), g(q_2),g(w_x),g(w_y))=(1,2,3,1,2,2,3,1)$, where w_x,w_y is the vertex in H adjacent to x and y, respectively. This completes the proof. ■

Lemma 3.8 *Let G be a triangle-free subcubic graph and H a subgraph of G. Let f be a proper $(G-H)$-rainbow-free 3-coloring of $G-H$. If H is isomorphic to R_0, then f can be extended to a proper G-rainbow-free 3-coloring g of G.*

Proof. It is clear that no graph in $\mathcal{L}_t$, for any t, contains R_0 as an induced subgraph as G is triangle-free. By Lemma 3.6, it is sufficient to show that f can be extend to a proper 3-coloring g of G such that H contains no rainbow L_0 with respect to g.

Let $V(H)=\{v_i,u_j,w:1\le i\le 4, j=1,2\}$, $(v_1,v_2,v_3,v_4,u_1,u_2)$ form a L_0, and w adjacent to u_1 and v_3. Note that if h is a proper 3-coloring of H such that $\{h(v_1),h(v_2),h(v_3)\}=\{1,2,3\}$, then H contains no rainbow L_0 with respect to h. Let $N_G(H)=\{x,y,z\}$ such that $xv_2,wy,u_2z\in E(G)$. Without loss of generality, we may assume that $f(z)=1$. Then we extend f to g by first defining $g(v_1)=1, g(v_2)\in\{2,3\}-\{f(x)\}, g(v_3)=\{2,3\}-\{g(v_2)\}$, and then it is easy to further define g on the remaining vertices such that g is a proper 3-coloring of G and G contains no rainbow L_0 with respect to g. ■

Lemma 3.9 *Let G be a K_4-free subcubic graph and S a subset of $E(G)$. If $G-S$ is $\{K_3,R_i:1\le i\le 7\}\cup\mathcal{L}'$-free, and no subgraph H of G isomorphic to a graph in $\mathcal{L}'$ satisfies that $H-S$ in $\mathcal{L}_t$ for some $t\ge 0$, then there is a proper $(G-S)$-rainbow-free 3-coloring f of G.*

Proof. We shall do induction on $|V(G)|$. If $G-S$ contains no L_0 as subgraph, then a proper $(G-S)$-rainbow-free 3-coloring exists by Brooks' Theorem. This proves the base case, and we assume that $G-S$ contains L_0 as a subgraph. Let H be a maximal subgraph of $G-S$ induced by a set T which is of the form $\bigcup_{i=1}^{k}H_i$ for some positive integer k, where each H_i is isomorphic to L_0. That is, if T' is a subset of $V(G)$ with $(G-S)[T']$ isomorphic to L_0, then $T'\cap T=\emptyset$ or $T'\subseteq T$. By Lemma 3.6, H is isomorphic to R_0 or a graph in $\mathcal{L}_t$ for some nonegative integer t. Note that if H is isomorphic to a graph in $\mathcal{L}_t$, then H is an induced subgraph of G, otherwise, the subgraph $G[V(H)]$ of G induced by $V(H)$ is isomorphic to a graph in $\mathcal{L}'$ and $G[V(H)]-S=H$ is in $\mathcal{L}_t$.

If H is isomorphic to R_0 and H is an induced subgraph of G, or H is in $\mathcal{L}_t$ such that $N_G(H)$ is not an independent set, or $\deg_G(v)=2$ for some $v\in V(H)$, then we can obtain a proper $(G-S)$-rainbow-free 3-coloring of G by extending the proper $(G'-S')$-rainbow-free

3-coloring of $G - V(H)$ obtained from applying induction to the graph $G' = G - V(H)$ and the set $S' = S \cap E(G - V(H))$, by Lemmas 3.8 and 3.7. Similarly, if H is isomorphic to R_0 but H not an induced subgraph of G, then $|N_G(H)| \leq 1$, and it is easy to extend the coloring obtained from applying induction on $G - V(H)$ and $S \cap E(G - V(H))$ to be a proper $(G - S)$-rainbow-free 3-coloring. So we may assume that H is in $\mathcal{L}_t$, $N_G(H)$ is an independent set, and $\deg_G(v) = 3$ for every $v \in V(H)$.

Assume that $|N_G(H)| \leq 3$, then some vertex x in $N_G(H)$ has degree at most one in $G - V(H)$. Let y be a vertex in $N_G(H) - \{x\}$, and let $G'' = (G - V(H)) + xy$ and $S'' = (S \cap E(G - V(H))) \cup \{xy\}$. So G'' is K_4-free, and no subgraph H' of G'' isomorphic to a graph in $\mathcal{L}'$ satisfies that $H' - S''$ is in $\mathcal{L}_t$ for some $t \geq 0$. Let f be the proper $(G'' - S'')$-rainbow-free 3-coloring of G'' obtained from applying induction to G'' and S''. Since $f(x) \neq f(y)$, we can extend f to a proper $(G - S)$-rainbow-free 3-coloring of G by Lemma 3.7. Hence, we assume that $|N_G(H)| = 4$.

It is clear that unless all vertices in $N_G(H)$ are in a subgraph of $G - H$ isomorphic to $L_2^{(r)}$ for some odd number r, there is a pair of vertices u, v in $N_G(H)$ such that $G - V(H) + uv$, denoted by G''', is a K_4-free subcubic graph and the edge set $(S \cap E(G - V(H))) \cup \{uv\}$, denoted by S''', has the property that $G''' - S'''$ is $\{K_3, R_i : 1 \leq i \leq 7\} \cup \mathcal{L}'$-free and no subgraph H' of G''' isomorphic to a graph in $\mathcal{L}'$ satisfies that $H' - S'''$ is in $\mathcal{L}_t$ for some $t \geq 0$, and we are done by extending the proper $(G''' - S''')$-rainbow-free 3-coloring obtained from applying induction to G''' and S''' in this case. Therefore, we assume that all vertices in $N_G(H)$ are in a subgraph J of $G - H$ isomorphic to $L_2^{(r)}$ for some odd number r. Note that there is a 4-cycle C such that $J - C$ is isomorphic to $L_2^{(r-1)}$ or L_0, and let a, c be the two diagonal vertices in C such that $a, c \in N_G(H)$. Apply induction to $(G - H - (J - C)) + ac$, denoted by G'''', and $(S \cap E(G - H - (J - C))) \cup \{ac\}$, denoted by S'''', to obtain a proper $(G'''' - S'''')$-rainbow-free 3-coloring of G'''', and then we can obtain a proper $(G - S)$-rainbow-free 3-coloring of G by first extending it to $G - H$, and then extending it to G. Since some vertex in $J - C$ has degree 2 in $G - H$, and a and c receive different colors, the above two procedures of extending colorings is allowed by Lemma 3.7. ■

Now, we are ready to prove Lemma 2.5.

Proof of Lemma 2.5: Given any proper 3-coloring f, we define $N(f)$ to be the number of vertices of degree three for which each of its three neighbors have the same color. Since G is a fractionally t-critical triangle-free subcubic graph with $t \geq 8/3$, G is $\{K_3, R_i : 1 \leq$

$i \le 7\} \cup \mathcal{L}'$-free by Lemma 3.3 and Lemma 3.5. Applying Lemma 3.9 by choosing $S = \emptyset$, there exists a proper G-rainbow-free 3-coloring of G. Let f be a proper G-rainbow-free 3-coloring of G with $N(f)$ as small as possible. We shall show that f is a good 3-coloring, and it is sufficient to show that every two color classes induce a good graph.

Claim 1: For every $1 \le i < j \le 3$, $G_{i,j}$ satisfies (G1).

Proof of Claim 1: Suppose to the contrary, and assume that $G_{2,3}$ does not satisfy (G1). Then there is a vertex x such that $f(x) = 2$, $f(z) = 3$ and $\deg_{G_{2,3}}(z) = 3$ for all $z \in N_{G_{2,3}}[x]$. However, replacing $f(x)$ by 1 reduces $N(f)$, and the coloring is still proper G-rainbow-free, a contradiction. □

Claim 2: For every $1 \le i < j \le 3$, $G_{i,j}$ satisfies (G1).

Proof of Claim 2: Suppose to the contrary, and assume that $G_{2,3}$ does not satisfy (G2). Then let x, y be two adjacent vertices such that $\deg_{G_{2,3}}(x) = \deg_{G_{2,3}}(y) = 3$ and $\deg_{G_{2,3}}(z) \ge 2$ for all $z \in N_{G_{2,3}}(x) \cup N_{G_{2,3}}(y)$, and $f(x) = 2$. Let x_1, x_2 be the two neighbors of x other than y, and y_1, y_2 be the two neighbors of y other than x. Let $x_{i,j}$ and $y_{i,j}$ be the neighbors of x_i and y_i other than x and y, repsectively, for $i = 1, 2$ and $j = 1, 2$. Let f_x (and f_y, respectively) be the coloring that is obtained from f by changing $f(x)$ (and $f(y)$, respectively) to 1. Notice that f_x and f_y are proper 3-colorings with $N(f_x), N(f_y) < N(f)$ by the degree condition of neighbors of x and y, so there exists a rainbow L_0 with respect to f_x and f_y. Furthermore, the rainbow L_0 with respect to f_x (or f_y, respectively) contains x (or y, respectively).

First, we assume that $x_{1,1} = x_{2,1}$. Then without loss of generality, we may assume that the rainbow L_0 with respect to f_x is formed by $(y, x, x_1, x_{1,2}, y_1, y_2)$. However, no f_y-rainbow L_0 exists in this case. Hence $N_{G_{2,3}}[x_1] \cap N_{G_{2,3}}[x_2] = \{x\}$ and $N_{G_{2,3}}[y_1] \cap N_{G_{2,3}}[y_2] = \{y\}$ by symmetry.

Second, we assume that $y_1 = x_{1,1}$ and $y_2 = x_{2,1}$. By the minimality of $N(f)$ and the triangle-freeness, $(x_{i,2}, x_i, x, x_{3-i}, x_{3-i,1}, x_{3-i,2})$ forms a rainbow L_0 with respect to f_x for some $i = 1, 2$. Then it is easy to check that there is no rainbow L_0 with respect to f_y. Similarly, if $y_1 = x_{1,1}$, $y_2 = x_{1,2}$, then either G has a cut-edge or $N(f) > N(f_x)$ and there is no f_x-rainbow L_0, a contradiction. Hence, $|\{x_{1,1}, x_{1,2}, x_{2,1}, x_{2,2}\} \cap \{y_1, y_2\}| \le 1$ by symmetry.

Third, we assume that $|\{x_{1,1}, x_{1,2}, x_{2,1}, x_{2,2}\} \cap \{y_1, y_2\}| = 1$, say $x_{2,1} = y_2$ and $y_{2,1} = x_2$. By symmetry, any f_x-rainbow L_0 is formed by $(x_{1,2}, x_1, x, x_2, x_{2,1}, x_{2,2})$, $(x_1, x, x_2, x_{2,2}, x_{1,1}, x_{1,2})$ or $(x_{1,2}, x_1, x, y, y_1, y_2)$ by the triangle-freeness. If $(x_{1,2}, x_1, x, x_2, x_{2,1}, x_{2,2})$ forms a f_x-rainbow L_0, then no f_y-rainbow L_0 exists. If $(x_1, x, x_2, x_{2,2}, x_{1,1}, x_{1,2})$ forms a rainbow L_0 with respect to f_x, then $x_{1,2} \neq y_1$ by the R_1-

freeness of G, and the unique rainbow L_0 with respect to f_y is formed by $(y_{2,2}, y_2, y, y_1, y_{1,1}, y_{1,2})$, where $y_{2,2}$ is the neighbor of y_2 other than x_2 and y. Let $f_{y_1,y_2,y_{2,2}}$ be the 3-coloring obtained from f by changing $(f(y_1), f(y_2), f(y_{2,2})) = (1, 1, 2)$. Then $f_{y_1,y_2,y_{2,2}}$ is a proper 3-coloring with $N(f_{y_1,y_2,y_{2,2}}) < N(f)$ without any rainbow L_0's. If $(x_{1,2}, x_1, x, y, y_1, y_2)$ forms a f_x-rainbow L_0, then no f_y-rainbow L_0 exists. Consequently, $\{x_{i,j} : i = 1, 2, j = 1, 2\}$ is disjoint from $\{y_1, y_2\}$.

Again by symmetry, we may assume that any f_x-rainbow L_0 is formed by $(x_{1,2}, x_1, x, x_2, x_{2,1}, x_{2,2})$ or $(x_{1,2}, x_1, x, y, y_1, y_2)$. For the latter case, no f_y-rainbow L_0 exists. So $(x_{1,2}, x_1, x, x_2, x_{2,1}, x_{2,2})$ forms a f_x-rainbow L_0, then we may assume that the f_y-rainbow L_0 is formed by $(y_{1,2}, y_1, y, y_2, y_{2,1}, y_{2,2})$ by symmetry. Let $f_{y_1,y_2,y_{1,2}}$ be the 3-coloring obtained from f by changing $(f(y_1), f(y_2), f(y_{1.2}))$ to $(1, 1, 2)$. Then $f_{y_1,y_2,y_{1,2}}$ is a proper 3-coloring of G with $N(f_{y_1,y_2,y_{1,2}}) \leq N(f)$, but no $f_{y_1,y_2,y_{1,2}}$-rainbow L_0 exists. So $N(f_{y_1,y_2,y_{1,2}}) = N(f)$ and $f(y_{1,1,1}) = f(y_{1,1,2}) = 1$, where $y_{1,1,1}$ and $y_{1,1,2}$ are the two neighbors of $y_{1,1}$ other than y_1. In other words, $\deg_{G_{2,3}}(y_1) = 2$ and $\deg_{G_{2,3}}(y_{1,1}) = 1$. Similarly, we can define $f_{x_1,x_2,x_{1,2}}$ by changing $(f(x_1), f(x_2), f(x_{1,2}))$ to $(1, 1, 3)$, and then we obtain that $\deg_{G_{2,3}}(x_1) = 2$ and $\deg_{G_{2,3}}(x_{1,1}) = 3$. So $G_{2,3}$ satisfies (G2), a contradiction. $\square$

Claim 3: For every $1 \leq i < j \leq 3$, $G_{i,j}$ satisfies (G3).

Proof of Claim 3: Suppose to the contrary, and assume that $G_{2,3}$ does not satisfy (G3). Let $x, y, z \in V(G_{2,3})$, $x_1 \in N_{G_{2,3}}(x) \setminus \{y\}$, $y' \in N_{G_{2,3}}(y) \setminus \{x, z\}$, $y'_1 \in N_{G_{2,3}}(y') - \{y\}$, $z_1 \in N_{G_{2,3}}(z) \setminus \{y\}$ such that xyz is a path in $G_{2,3}$, $\deg_{G_{2,3}}(x) = \deg_{G_{2,3}}(y) = \deg_{G_{2,3}}(z) = 3$, $\deg_{G_{2,3}}(x_1) = \deg_{G_{2,3}}(z_1) = 1$, $\deg_{G_{2,3}}(y') = 2$, and $\deg_{G_{2,3}}(y'_1) \geq 2$, and $f(y) = 2$. Let r_y be the 3-coloring obtained from f by changing $f(y)$ to 1. Since $N(r_y) < N(f)$, there is a rainbow L_0 with respect to r_y. Without loss of generality, we may assume that the r_y-rainbow L_0 is formed by $(z, y, y', z_{1,1}, z_1, z_2)$, where $N_{G_{2,3}}(z) = \{z_1, z_2, y\}$ and $N_G(z_1) = \{z, z_{1,1}, z_{1,2}\}$. Let $r_{z,y',z_{1,1}}$ be the 3-coloring obtained from f by changing $(f(z), f(y'), f(z_{1,1}))$ to $(1, 1, 3)$. We derive a contradiction since no $r_{z,y',z_{1,1}}$-rainbow L_0 exists and $N(r_{z,y',z_{1,1}}) < N(f)$. $\square$

Claim 4: For every $1 \leq i < j \leq 3$, $G_{i,j}$ satisfies (G4).

Proof of Claim 4: Suppose to the contrary, and assume that $G_{2,3}$ does not satisfy (G4). Let $x, y, z \in V(G_{2,3})$ such that xyz is a path in $G_{2,3}$, $\deg_{G_{2,3}}(x) = \deg_{G_{2,3}}(y) = \deg_{G_{2,3}}(z) = 3$, $\deg_{G_{2,3}}(z_1) \geq 2$, and $\deg_{G_{2,3}}(z_2) \geq 2$, where $N_G(z) = \{z_1, z_2, y\}$, and $f(z) = 2$, but none of z_1, z_2 is a support vertex of degree two in $G_{2,3}$. Let p_z be the 3-coloring obtained from f by changing $f(z)$ to 1. Since $\deg_{G_{2,3}}(z_1) \geq 2$ and $\deg_{G_{2,3}}(z_2) \geq 2$, we have that $N(p_z) < N(f)$, so there is a rainbow L_0 with respect to p_z. Without loss of generality, we may assume that the rainbow L_0 is formed by $(z_1, z, z_2, z_{2,1}, z_{1,1}, z_{1,2})$, where $N_{G_{2,3}}(z_1) = \{z, z_{1,1}, z_{1,2}\}$ and $N_{G_{2,3}}(z_2) = \{z, z_{2,1}, z_{2,2}\}$. Notice that it implies that $\deg_{G_{2,3}}(z_1) = 3$ and $\deg_{G_{2,3}}(z_2) = 2$. Let $p_{z_1,z_2,z_{2,1}}$ be the 3-coloring obtained from f by changing $(f(z_1), f(z_2), f(z_{2,1}))$ to $(1, 1, 3)$. Since there is no rainbow L_0 with respect to $p_{z_1,z_2,z_{2,1}}$, we have $N(p_{z_1,z_2,z_{2,1}}) = N(f)$, and this implies that $\deg_{G_{2,3}}(z_{2,2}) = 1$. That is, z_2 is a suppoer vertex of degree two in $G_{2,3}$, a contradiction. $\square$

Together with Claims 1 thorugh 4, f is a good 3-coloring. This completes the proof of Lemma 2.5. $\square$

4 Avoiding coloring

The purpose of this section is to obtain lemmas that will be used in Section 5.

Recall that $f : V(G) \to 2^{[14]}$ is an *F-avoiding coloring* of a graph G if $f(v)$ is disjoint from $F(v) \cup f(u)$ for every two adjacent vertices u and v, where $F : V(G) \to 2^{[14]}$.

Lemma 4.1 *Let $0 \leq r_1 \leq 2$ and $0 \leq r_{2k+1} \leq 2$, P an odd path $v_1v_2...v_{2k+1}$, where $k \geq 1$, and $F : V(P) \to 2^{[14]}$ a function with $|F(v_j)| \leq 3 + r_j$ for $j = 1, 2k+1$ and $|F(v_i)| = 2$ for all $2 \leq i \leq 2k$, and $|F(v_1) \cap F(v_2)| \leq r_1$, $|F(v_{2k}) \cap F(v_{2k+1})| \leq r_{2k+1}$. Then there is an F-avoiding coloring f of P such that $|f(v_j)| = 6$ for every $2 \leq j \leq 2k$, and $|f(v_j)| = 8 - r_j$ for $j = 1, 2k+1$.*

Proof. We shall prove this lemma by doing induction on k. Note that we may assume that $|F(v_j)| = 3 + r_j$ for $j = 1, 2k+1$. Let S_1 be a subset of $F(v_1) - F(v_2)$ of size 3, and S_{2k+1} be a subset of $F(v_{2k+1}) - F(v_{2k})$ of size 3. When $k = 1$, define $f : V(P) \to 2^{[14]}$ by letting $f(v_2) = S_1 \cup S_{2k+1} \cup S$, where S is a subset of $[14] - (S_1 \cup S_{2k+1} \cup F(v_2))$ of size $6 - |S_1 \cup S_{2k+1}|$, and letting $f(v_1)$ and $f(v_{2k+1})$ be a subset of $[14] - (F_1 \cup f(v_2))$ of size $8 - r_1$ and a subset of $[14] - (F_{2k+1} \cup f(v_{2k}))$ of size $8 - r_{2k+1}$, respectively. Notice that $|F_1 \cup f(v_2)| \leq 6 + r_1$ and $|F_{2k+1} \cup f(v_{2k})| \leq 6 + r_{2k+1}$, so $f(v_1)$ and $f(v_{2k+1})$ are well-defined. This proves the base case.

Now, we assume that $2k + 1 \geq 5$ and the lemma holds for all odd path of order at least three but less than $2k+1$. Define $F' : \{v_i : 3 \leq i \leq 2k+1\} \to 2^{[14]}$ by $F'(v_3) = F(v_3) \cup S_1$ and $F'(v_i) = F(v_i)$ for all $3 < i \leq 2k+1$. Hence, by the induction hypothesis, there is an F'-avoiding coloring f' of $P - \{v_1, v_2\}$ such that $|f'(v_{2k+1})| = 8 - r_{2k+1}$ and $|f'(v_i)| = 6$ for all $3 \leq i \leq 2k+1$. Then define $f : V(P) \to 2^{[14]}$ by $f(v_i) = f'(v_i)$ for all $3 \leq i \leq 2k+1$, $f(v_2) = S_1 \cup T_2$, where T_2 is a subset of $[14] - (F(v_2) \cup f'(v_3) \cup S_1)$ of size 3, and $f(v_1)$ be a subset of $[14]-(F_1\cup f(v_2))$ of size $8-r_1$. Notice that f is an F-avoiding coloring of P, so this proves the lemma. ■

Lemma 4.2 *Let* P *be an even path* $v_1v_2...v_{2k}$, *where* $k\geq 1$, *and* $F: V(P)\rightarrow 2^{[14]}$ *a function with* $|F(v_j)|\leq 4$ *for* $j=1,2k$ *and* $|F(v_i)|=2$ *for all* $2\leq i\leq 2k-1$. *Then there is an* F*-avoiding coloring* f *of* P *such that* $|f(v_j)|=8-\deg_P(v_j)$ *for* $1\leq j\leq 2k$ *if one of the following holds:*

1. $k=1$ *and* $F(v_1)\cap F(v_2)=\emptyset$*;*
2. $k=2$, $F(v_1)\cap F(v_2)=\emptyset$, $F(v_3)\cap F(v_4)=\emptyset$, *and* $|F(v_1)\cap F(v_4)|\leq 2$*;*
3. $k\geq 3$, $F(v_{2i+4})\subseteq F(v_{2i+1})$ *for* $0\leq i\leq k-3$, $F(v_{2k-3})\subseteq F(v_{2k})$, $(F(v_1)-F(v_4))\cap(F(v_{2k})-F(v_{2k-3}))=\emptyset$, $F(v_1)\cap F(v_2)=\emptyset$, $F(v_{2j-1})\cap F(v_{2j})=\emptyset$ *for* $2\leq j\leq k-1$, *and* $F(v_{2k-1})\cap F(v_{2k})=\emptyset$*;*
4. $k\geq 3$, $F(v_1)\cap F(v_2)=\emptyset$, $F(v_1)\cap F(v_4)=\emptyset$, $F(v_3)\cap F(v_4)=\emptyset$, *and* $F(v_{2k-1})\cap F(v_{2k})=\emptyset$*;*
5. $k\geq 3$, $F(v_{2i+4})\subseteq F(v_{2i+1})$ *for* $0\leq i\leq k-3$, $F(v_1)\cap F(v_2)=\emptyset$, $f(v_{2j-1})\cap f(v_{2j})=\emptyset$ *for* $2\leq j\leq k-2$, $F(v_{2k-3})\cap F(v_{2k})=\emptyset$, *and* $F(v_{2k-1})\cap F(v_{2k})=\emptyset$*;*
6. $k\geq 4$, $F(v_1)\cap F(v_2)=\emptyset$, *and there exists* $0\leq s\leq k-4$ *such that* $F(v_{2i+4})\subseteq F(v_{2i+1})$ *for* $0\leq i\leq s$, $F(v_{2i+3})\cap F(v_{2i+4})=\emptyset$ *for* $0\leq i\leq s-1$, $F(v_{2s+3})\cap F(v_{2s+6})=\emptyset$, $F(v_{2s+5})\cap F(v_{2s+6})=\emptyset$, $(F(v_1)-F(v_4))\cap F(v_{2s+6})=\emptyset$, *and* $F(v_{2k-1})\cap F(v_{2k})=\emptyset$*;*
7. $k\geq 2$, $|F(v_1)|=2$, $|F(v_1)\cap F(v_2)|\leq 1$, *and* $F(v_{2k-1})\cap F(v_{2k})=\emptyset$*;*
8. $k\geq 3$, $F(v_3)\cup F(v_4)\subseteq F(v_1)$, $F(v_1)\cap F(v_2)=\emptyset$, *and* $F(v_{2k-1})\cap F(v_{2k})=\emptyset$*;*
9. $k=4$, $F(v_1)=F(v_4)\cup F(v_6)$, $F(v_1)\cap F(v_2)=\emptyset$, $(F(v_5)\cup F(v_7))\cap F(v_8)=\emptyset$*;*
10. $k\geq 5$, $F(v_1)=F(v_4)\cup F(v_6)$, *and* $F(v_1)\cap F(v_2)=(F(v_3)\cup F(v_5)\cup F(v_7))\cap F(v_8)=F(v_{2k-1})\cap F(v_{2k})=\emptyset$*;*
11. $k\geq 4$, $F(v_1)=F(v_4)\cup F(v_6)$, *and there exists* $0\leq t\leq k-4$ *such that* $F(v_{2i+5})\subseteq F(v_{2i+8})$ *for* $0\leq i\leq t$, $F(v_1)\cap F(v_2)=F(v_5)\cap f(V_6)=F(v_{2k-1})\cap F(v_{2k})=\emptyset$, $F(v_{2j+7})\cap F(v_{2j+8})=\emptyset$ *for* $0\leq j\leq t-1$, *and one of the following holds:*
    - *(a)* $t=k-4$, *and* $F(v_3)\cap(F(v_{2k})-F(v_{2k-3}))=\emptyset$.
    - *(b)* $k\geq 5$, $t=k-5$, $F(v_{2k-3})\cap F(v_{2k})=\emptyset$.
    - *(c)* $k\geq 6$, $t\leq k-6$, *and* $(F(v_3)\cup F(v_{2t+7})\cup F(v_{2t+9}))\cap F(v_{2t+10})=\emptyset$.

12. *$k \geq 2$, $|F(v_i)| \leq 4 - 2\deg_P(v_i)$ for some $1 \leq i \leq 2k$, $i \neq 2, 2k-1$, and $F(v_1) \cap F(v_2) = F(v_{2k-1}) \cap F(v_{2k}) = \emptyset$.*

Proof.

1. Statement 1 is clear by observating that there is a partition $X \cup Y$ of $[14]$ such that $|X| = |Y| = 7$, $X \cap F(v_1) = \emptyset$, and $Y \cap F(v_2) = \emptyset$.
2. We may assume that $|F(v_1)| = |F(v_4)| = 4$. Since $|F(v_1) \cap F(v_4)| \leq 2$, there are two disjoint sets X and Y with sizes 3, where $X \subseteq F(v_1)$ and $Y \subseteq F(v_4)$. Pick a subset Y' of $[14] - (X \cup Y \cup F(v_3))$ with size 3, and then assign $f(v_3) = Y \cup Y'$. Define $f(v_2)$ to be a subset of $[14] - (f(v_3) \cup F(v_2))$ containing X with size 6. Finally, pick a subset W of $[14] - (f(v_2) \cup F(v_1))$ with size 7. Notice that $|[14] - (f(v_2) \cup F(v_1))| \geq 7$ since $f(v_2) \cap F(v_1)$ contains X which has size 3. Similarly, there is a subset Z of $[14] - (f(v_3) \cup F(v_4))$ with size 7. Then assigning $f(v_1) = W$ and $f(v_4) = Z$ gives a desired F-avoiding coloring.
3. We shall do induction on k. Without loss of generality, we can assume that $F(v_1)$ has size four. Let S be a subset of $F(v_1)$ containing $F(v_1) - F(v_4)$ with size 3. Define $F' : \{v_i : 3 \leq i \leq 2k\} \to 2^{[14]}$ by letting $F'(v_3) = F(v_3) \cup (F(v_1) - F(v_4))$, and $F'(v_i) = F(v_i)$ for all $i > 3$. Then there is an F'-avoiding coloring f' of the subpath $v_3 v_4 ... v_{2k}$ by statement 2 (when $k = 3$) and by the induction hypothesis (when $k > 3$). Define $f(v_i) = f'(v_i)$ for $i \geq 4$, and define $f(v_3)$ to be a subset of $f'(v_3) - S$ with size 6, $f(v_2)$ to be a subset of $[14] - (F(v_2) \cup f(v_3))$ containing S with size 6, and $f(v_1)$ to be a subset of $[14] - (f(v_2) \cup F(v_1))$ with size 7. Note that $|f'(v_3) \cap S| \leq 1$, so f is well-defined.
4. We may assume that $|F(v_1)| = |F(v_{2k})| = 4$. Let S be a subset of $[14] - (F(v_1) \cup F(v_3) \cup F(v_4))$ of size 3. Define $F' : \{v_i : 4 \leq i \leq 2k\} \to 2^{[14]}$ by $F'(v_4) = F(v_4) \cup S$, and $F'(v_i) = F(v_i)$ for all $i > 4$. By Lemma 4.1, there is an F'-avoiding coloring f' with $|f'(v_{2k})| = 7$ and $|f'(v_i)| = 6$ for all $4 \leq i < 2k$. Pick a subset S' of $[14] - (f'(v_4) \cup F(v_3) \cup F(v_4) \cup S)$ with size 1. Assign $f(v_3) = F(v_4) \cup S \cup S'$ and $f(v_i) = f'(v_i)$ for all $i \leq 4 \leq 2k$. Let T be a subset of $[14] - (F(v_2) \cup f(v_3))$ with size $6 - |F(v_1) - S'|$, and then assign $f(v_2) = (F(v_1) - S') \cup T$. Finally, assign $f(v_1)$ by any subset of $[14] - (F(v_1) \cup f(v_2))$ with size 7. It is clear that f is a desired F-avoiding coloring.
5. We shall do indeuction on k. We may assume that $|F(v_1)| = 4$. Assume that $k = 3$. If $F(v_3) = F(v_4)$, then define $S = F(v_4) \cup \{s\}$, where $s \in F(v_1) - F(v_4)$, and $S' = \{s\}$; otherwise, let $S = (F(v_1) - F(v_4)) \cup \{s\}$, where $s \in F(v_4)$ such that $|S \cap F(v_3)|$

is as large as possible, and let S' be a subset of $(S \cup F(v_3)) - F(v_4)$ with size $|S \cup F(v_3)| - 2$ such that $|S' \cap F(v_3)|$ is as large as possible. In fact, $|S' \cap F(v_3)| \geq |S'| - 1$. Then there is a subset X of $F(v_6)$ of size $|F(v_6)| - 1$ such that $X \cap S' = \emptyset$. Define an F-avoiding coloring $f : V(G) \to 2^{[14]}$ by assigning $f(v_5)$ any subset of $[14] - (F(v_5) \cup S')$ containing X with size 6, $f(v_6)$ any subset of $[14] - (f(v_5) \cup F(v_6))$ with size 7, $f(v_4)$ any subset of $[14] - (F(v_4) \cup f(v_5))$ containing S' with size 6, $f(v_3)$ any subset of $[14] - (F(v_3) \cup f(v_4) \cup S)$ with size 6, $f(v_2)$ any subset of $[14] - (F(v_2) \cup f(v_3))$ containing S with size 6, and $f(v_1)$ any subset of $[14] - (F(v_1) \cup f(v_2))$ with size 7. This prove the base case of the induction.

When $k > 3$, let S_2 be a subset of $F(v_1) - F(v_2)$ with size 3 such that $|S_2 \cap F(v_4)| \leq 1$. Define $F' : \{v_i : 3 \leq i \leq 2k\} \to 2^{[14]}$ by letting $F'(v_3)$ be a subset of $(F(v_3) \cup S_2) - F(v_4)$ with size $|F(v_3) \cup S_2| - 1$, and $F'(v_i) = F(v_i)$ for $i \geq 4$. By induction, there is an F'-avoiding coloring f' of the subspath $v_3 v_4 ... v_{2k}$ such that $|f'(v_3)| = |f'(v_{2k})| = 7$ and $|f'(v_i)| = 6$ for $4 \leq i \leq 2k - 1$. And define $f(v_i) = f'(v_i)$ for $i \geq 4$, $f(v_3)$ to be a subset of $f'(v_3) - S_2$ with size 6, and $f(v_2)$ to be a superset of S_2 disjoint from $f(v_3) \cup F(v_2)$ with size 6, and $f(v_1)$ to be a subset of $[14] - (f(v_2) \cup F(v_1))$ with size 7. It is easy to check that f is desired.

6. Induction on k. Assume that $|F(v)| = 6 - 2\deg_G(v)$ for all v. When $k = 4$, $s = 0$, and let S_5 be a superset of $F(v_6)$ disjoint from $F(v_1) \cup F(v_3) \cup F(v_5)$ with size 5. By Lemma 4.1, there is an F-avoiding coloring f of the subpath $v_6 v_7 ... v_{2k}$ such that $f(v_6) \cap S_5 = \emptyset$ and $|f(v_i)| = 8 - \deg_P(v_i)$ for $6 \leq i \leq 2k$. Now, we extend f to $V(G)$. First, define $f(v_5)$ be a superset of S_5 disjoint from $F(v_5) \cup f(v_6)$ with size 6. If $F(v_3) \neq F(v_4)$, then let S_2 be a subset of $F(v_1) - F(v_2)$ with size 3 such that $|(S_2 \cup F(v_3)) \cap F(v_4)| \leq 1$, and let S_4 be a subset of $S_2 \cup F(v_3) - (F(v_4) \cup f(v_5))$ with size $|S_2 \cup F(v_3)| - 2$. If $F(v_3) = F(v_4)$, then let S_2 be a subset of $F(v_1) - F(v_2)$ containing $F(v_4)$ with size 3 such that $(S_2 - F(v_4)) \cap f(v_5) = \emptyset$, and let S_4 be a subset of $S_2 - f(v_5)$ with size 1. Define $f(v_4)$ to be a superset of S_4 disjoint from $F(v_4) \cup f(v_5)$ with size 6, $f(v_3)$ to be a subset of $[14] - (f(v_4) \cup F(v_3))$ with size 6, $f(v_2)$ to be a superset of S_2 disjoint from $F(v_2) \cup f(v_3)$ with size 6, and $f(v_1)$ to be a subset of $[14] - (F(v_1) \cup f(v_2))$ with size 7. This proves the base case.

   Now assume that the statement holds for every smaller $k \geq 4$. Let S be a subset of $F(v_1) - F(v_2)$ containing $F(v_1) - F(v_4)$ with size 3. Define $F' : \{v_i : 3 \leq i \leq 2k\} \to 2^{[14]}$ by letting $F'(v_3)$ be $(F(v_1) - F(v_4)) \cup F(v_3)$, and $F'(v) = F(v)$ for every other v.

By induction, there is an F'-avoiding coloring f' of the subpath $v_3v_4...v_{2k}$. Extend f' to f by defining $f(v_i) = f'(v_i)$ for $i \geq 4$, $f(v_3)$ to be a subset of $f'(v_3) - S$ with size 6, and $f(v_2)$ to be a superset of S disjoint from $F(v_2) \cup f(v_3)$ with size 6, and $f(v_1)$ to be a subset of $[14] - F(v_1) - f(v_2)$ with size 7.

7. Induction on k. The case that $k = 2$ is easy. When $k \geq 3$, let S be a subset of $F(v_1) - F(v_2)$ with size 1. Define $F' : \{v_i : 3 \leq i \leq 2k\} \to 2^{[14]}$ by letting $F'(v_3)$ be a subset of $S \cup F(v_3)$ with size $|S \cup F(v_3)| - 1$ such that $|F'(v_3) \cap F(v_4)| \leq 1$, and define $F'(v_i) = F(v_i)$ for $i \geq 4$. By induction, there is an F'-avoiding coloring f' of the subpath. Extend f' to f by defining $f(v_i) = f'(v_i)$ for $4 \leq i \leq 2k$, $f(v_3)$ to be a subset of $f'(v_3) - S$ with size 6, $f(v_2)$ to be a superset of S disjoint from $F(v_2) \cap f(v_3)$ with size 6, and define $f(v_1)$ to be a subset of $[14] - (f(v_2) \cup F(v_1))$ with size 7.

8. Let S be a subset of $F(v_1) - F(v_2)$ containing $F(v_3)$ with size 3. Define $F' : \{v_i : 3 \leq i \leq 2k\} \to 2^{[14]}$ by letting $F'(v_3)$ be a subset of S with size 2 such that $|F'(v_3) \cap F(v_4)| \leq 1$, and define $F'(v_i) = F(v_i)$ for $i \geq 4$. By statement 7, there is an F'-avoiding coloring f' of the subpath. And it is easy to use the same argument as in the proof of statement 7 to extend f' to an F-avoiding coloring f of P.

9. Let S_2 be a subset of $F(v_1) - F(v_2)$ containing $F(v_6)$ with size $|F(v_1)| - 1$, S_4 be a subset of $(S_2 \cup F(v_3)) - F(v_4)$ with size $|S_2 \cup F(v_3)| - 2$, S_6 be s subset of $(S_4 \cup F(v_5)) - F(v_6)$ with size $|S_4 \cup F(v_5)| - 2$ such that $|S_6 \cap F(v_5)|$ is as large as possible. It is not hard to see that $|S_6 \cap F(v_8)| \leq 1$, so there is a subset S_7 of $F(v_8) - (F(v_7) \cup S_6)$ with size $|F(v_8)| - 1$. Therefore, we can define a desired F-avoiding coloring f of P by defining $f(v_7)$ to be a superset of S_7, and then defining $f(v_8)$ to be a subset of $[14] - (F(v_8) \cup f(v_7))$ with size 7, and $f(v_j)$ from $j = 6$ down to 1 to be a set such that $f(v_j)$ is disjoint from $F(v_j) \cup f(v_{j+1})$ and contains S_j whenever S_j is defined.

10. We may assume that $F(v_3) \neq F(v_4)$ by statement 8. Let S_7 be a subset of $[14] - (F(v_3) \cup F(v_5) \cup F(v_7))$ containing $F(v_8)$ with size 5 such that $|S_7 \cap F(v_4)|$ is as small as possible. By Lemma 4.1, there is an F-avoiding coloring f' of the subpath $v_8v_9...v_{2k}$ such that $|f'(v_i)| = 6$ for $8 \leq i \leq 2k-1$ and $|f'(v_{2k})| = 7$. Define $f(v_i) = f'(v_i)$ for $8 \leq i \leq 2k$, and $f(v_7)$ to be a superset of S_7 with size 6 disjoint from $F(v_7) \cup f(v_8)$. Let S_5 be the subset of $f(v_7) \cup F(v_6) - F(v_5)$ with size $|f(v_7) \cup F(v_6)| - 2$ such that $|S_5 \cap F(v_6)| \leq 1$, and S_3 be the subset of $S_5 \cup F(v_4) - F(v_3)$

with size $|S_5 \cup F(v_4)| - 2$ such that $|S_3 \cap F(v_1)| = |S_3 \cap (F(v_4) \cup F(v_6))| \leq 1$. Define $f(v_2)$ to be a subset of $[14] - (F(v_2) \cup S_3)$ with size 6 such that $|f(v_2) \cap F(v_1)| \geq |F(v_1)| - 1$. And then define $f(v_1)$ to be a subset of $[14] - (F(v_1) \cup f(v_2))$ with size 7, and define$f(v_j)$ from $j = 3$ through 6 such that $f(v_j)$ is disjoint from $F(v_j) \cup f(v_{j-1})$ and contains S_j whenever S_j is defined.

11. Define S_2 to be a subset of $F(v_1)$ with size $|F(v_1)| - 1$ such that $|S_2 \cap F(v_3)|$ is as large as possible, and define S_4 to be a subset of $(S_2 \cup F(v_3)) - F(v_4)$ with size $|S_2 \cup F(v_3)| - 2$ such that $|S_4 \cap F(v_3)|$ is as large as possible. Note that $|S_4 \cap F(v_6)| \leq 1$. Define $F' : \{v_i : 5 \leq i \leq 2k\} \to 2^{[14]}$ by letting $F'(v_5)$ be a subset of $(S_4 \cup F(v_5)) - F(v_6)$ with size $|S_4 \cup F(v_5)| - 1$, and $F'(v_i) = F(v_i)$ for $6 \leq i \leq 2k$. By statements 2, 3, 5, and 6, there is an F'-avoiding coloring f' of the subpath $v_5v_6...v_{2k}$ such that $|f'(v_5)| = |f'(v_{2k})| = 7$ and $|f'(v_i)| = 6$ for $6 \leq i \leq 2k - 1$. Define an F-avoiding coloring f of P by letting $f'(v_i) = f(v_i)$ for $6 \leq i \leq 2k$, and assigning $f(v_5)$ a subset of $f'(v_5) - S_4$ with size 6, and from $j = 4$ down to 1, define $f(v_j)$ to be a subset of $[14] - (F(v_j) \cup f(v_{j+1}))$ containing S_j (if S_j is defined) with size 6 (size 7 when $j = 1$). It is clear that f is a desired F-avoiding coloring of P.

12. Without loss of generality, we may assume that i is odd, so $i \leq 2k - 3$. Define S_2 to be a subset of $F(v_1) - F(v_2)$ with size $|F(v_1)| - 1$, and define S_{2j} to be a subset of $(S_{2j-2} \cup F(v_{2j-1})) - F(v_{2j})$ with size $|S_{2j-2} \cup F(v_{2j-1})| - 2$ for $2 \leq j \leq k - 1$. Since $i \leq 2k - 3$, $|S_{2k-2}| \leq 1$. Define $f(v_{2k-1})$ to be a subset of $[14] - (S_{2k-2} \cup F(v_{2k-1}))$ with size 6 such that $|f(v_{2k-1}) \cap F(v_{2k})| \geq |F(v_{2k})| - 1$, and then we define $f(v_i)$ for $i \neq 2k - 1$ containing S_i (if S_i is defined) to make f a desired F-avoiding coloring.

This completes the proof. ■

Recall that for every even numbers $a, b \geq 4$, $H_{a,b}$ is the graph obtained from two disjoint paths P_a and P_b by adding an edge incident with one support vertex in P_a and one support vertex in P_b.

Lemma 4.3 *Let $a, b \geq 4$ be two even numbers. Let $V(H_{a,b}) = \{v_i, u_j : 1 \leq i \leq a, 1 \leq j \leq b\}$ and $E(H_{a,b}) = \{v_iv_{i+1}, u_ju_{j+1}, v_2u_2 : 1 \leq i \leq a - 1, 1 \leq j \leq b - 1\}$. And let $F : V(H_{a,b}) \to 2^{[14]}$ with $|F(v)| \leq 6 - 2\deg_{H_{i,j}}(v)$ for every vertex v in $H_{i,j}$. If $F(u_{b-1}) \cap F(u_b) = \emptyset$, and any condition in Lemma 4.2 holds for the path $v_1v_2...v_a$ or the path $v'_1v'_2...v'_a$, where $v'_i = v_{a+1-i}$. then there is an F-avoiding coloring f such that $|f(v)| = 8 - \deg_{H_{a,b}}(v) - 1_{\{u_2\}}(v)$.*

Proof. We may assume that $F(u_1) \neq \emptyset$. Let S_2 be a subset of $F(u_1)$ with size 1. Apply Lemma 4.2, there is an F-avoiding coloring f' for the path $v_1 v_2 ... v_a$ such that $|f'(v_2)| = 6$ and $|f'(v_i)| = 8 - \deg_{H_{a,b}}(v_i)$ for every $i \neq 2$. Define $f(v_2)$ to be a subset of $f'(v_2) - S_2$ with size 5, and $f(v_i) = f'(v_i)$ for every other i. Let S_3 be a subset of $f(v_2) - (S_2 \cup F(u_3))$ with size 1. For every j with $4 \leq j \leq b-2$, define S_j to be a subset of $(S_{j-2} \cup F(u_{j-1})) - F(u_j)$ with size $|S_{j-2} \cup F(u_{j-1})| - 2$. Define $f(u_{b-1})$ to be a superset of $(S_{b-3} \cup F(u_{b-2}) \cup F(v_b)) - (S_{b-2} \cup F(v_{b-1}))$ with size at 6 such that $|f(v_{b-1}) \cap (S_{b-3} \cup F(u_{b-2}))| \geq |S_{b-3} \cup F(u_{b-2})| - 2$ and $|f(v_{b-1}) \cap F(v_b)| \geq |F(v_b)| - 1$. Then, for $j = b-2$ down to 3 it is easy to define $f(u_j)$ to be a superset of S_j disjoint from $F(u_j) \cup f(u_{j+1}) \cup S_{j-1}$ such that $|f(u_j)| = 6$, and define $f(u_b)$ to be a subset of $[14] - F(v_b) - f(v_{b-1})$ with size 7. Define $f(u_2)$ to be a subset of $[14] - (f(v_2) \cup f(u_3))$ containing S_2 with size 4. Note that such set exists since $f(v_2) \cap f(u_3) \neq \emptyset$, which implies that $|f(v_2) \cup f(u_3)| \leq 10$. Finally, define $f(u_1)$ to be a subset of $[14] - f(u_2)$ with size 7. So it is clear that f is an F-avoiding coloring such that $|f(v)| = 8 - \deg_{H_{a,b}}(v) - 1_{\{u_2\}}(v)$ for every vertex v. ■

A *kernel* of a digraph D is an independent set K of D such that $N_D^+(u) \cap K \neq \emptyset$ for every vertex $u \notin K$. Richardson [12] proved that every diagraph having no odd directed cycle has a nonempty kernel. Now we generalize a theorem given by Gutner and Tarsi [5] as follows.

Lemma 4.4 *Let $D = (V, E)$ be a digraph, and let $r_D : V \to \mathbb{N} \cup \{0\}$ be a function. For each $v \in V$, let $S(v)$ be a set of size at least $r_D(v) + \sum_{u \in N_D^+(v)} r_D(u)$. If D contains no odd directed cycle, then there exist subsets $C(v) \subseteq S(v)$ with size $r_D(v)$ for all $v \in V$ such that $C(u) \cap C(v) = \emptyset$ for every two adjacent verteces u, v of D.*

Proof. We shall do induction on $|V|$ and $\sum_{v \in V} r_D(v)$. When $|V| = 1$, the lemma is obviouly true no matter what $\sum_{v \in V} r_D(v)$ is. Assume that $|V| > 1$, and the lemma holds for all digraph D' without odd directed cycles with order less than D and any nonnegative integral function $r_{D'}$ defined on $V(D')$, and also holds for D' with order same as D but $\sum_{v \in V(D')} r_{D'}(v)$ less than $\sum_{v \in V} r_D(v)$.

We may assume that $r_D(v) > 0$ for every vertex v in D, otherwise, we can just remove v from D and apply induction. Pick an element c in $S(v)$ for some vertex v, let H be the subdigraph of D induced by vertices u with $c \in S(u)$. H does not contain odd directed cycles as D does not, so there is a nonempty kernel K of H. Note that if $S(v) = \{c\}$, then $r_D(v) = 1$ and $N_D^+(v) = \emptyset$, so $v \in K$. Define $S'(u) = S(u) - \{c\}$ if $c \in S(u)$, and define $S'(u) = S(u)$ if $c \notin S(u)$. Define $r'_D(v) = r_D(v) - 1$ if $v \in V \cap K$, and $r'_D(v) = r_D(v)$ if $v \in V - K$. Note that if $v \in V \cap K$, then $|S'(v)| = |S(v)| - 1 \geq r_D(v) + \sum_{u \in N_D^+(v)} r_D(u) - 1 = r'_D(v) + \sum_{u \in N_D^+(v)} r'_D(u)$. Also, since every vertex not in K has an out-neighbor in K, $|S'(v)| \geq r'_D(v) + \sum_{u \in N_D^+(v)} r'_D(u)$ for vertex v not in K. Hence, by the induction hypothesis, there are subsets $C'(v)$ of $S'(v)$ for every vertex v of D' such that $|C'(v)| = r'_D(v)$ and $C'(u) \cap C'(v) = \emptyset$ for every two adjacent vertices u and v. Now define $C(v) = C'(v) \cup \{c\}$ if $v \in V \cap K$, and $C(v) = C'(v)$ if $v \in V - K$. Then $|C(v)| \geq r_D(v)$ for every vertex v of D, and $C(u) \cap C(v) = \emptyset$ for every two adjacent vertices u, v. This proves the lemma. ■

Recall that every fractionally t-critical triangle-free subcubic graph, where $t \geq 8/3$, has a proper 3-coloring of G such that every two color classes induce a good graph by Lemma 2.5.

Lemma 4.5 *Let G be a good subcubic graph. If X is a connected subgraph of order at least two, then $G - X$ is good.*

Proof. It is clear that $G - X$ satisfies (G1), (G2) and (G4) since G satisfies them. Furthermore, $G - X$ satisfies (G3) since G satisfies (G1), (G2) and (G4). ■

Given a matching M (not necessary induced) of G such that M does not saturate any support vertex, we call a function $F : V(G) \to 2^{[14]}$ *obeys M* if the following hold:

1. $|F(v)| \leq 6 - 2\deg_G(v)$ for every vertex v in G,
2. $F(v)$ is disjoint from $F(u)$ for every leaf v in G and the neighbor u of v,
3. $|F(x) \cap F(y)| \leq 1$ for every $xy \in M$.

We say that G is *(M, I)-tractable* if M is a matching of G saturating no support vertices, and I is an independent set of G such that every vertex in I has degree three, and for every F that obeys M, there exist two F-avoiding colorings f_1 and $f_2 : V(G) \to 2^{[14]}$ of G, where f_1, f_2 might not be distinct, such that $|f_1(v)| + |f_2(v)| = 16 - 2\deg_G(v) - 2 \cdot 1_I(v)$. In this case, we say (f_1, f_2) is a *F-avoiding pair* of G with respect to I. Note that every path of odd order is $(\emptyset, \emptyset)$-tractable by Lemma 4.1.

Lemma 4.6 *Let G be a connected subcubic graph which is not a path, and let u be a support vertex of degree two and $N_G(u) = \{v, w\}$, where v is a leaf and w is not a support vertex. Let $G' = G - \{u, v\}$. If G' is (M', I')-tractable for some matching M' of G' and independent set I' of G', then G is (M, I')-tractable for some mathching M of G.*

Proof. If w is a leaf in G' or w is saturated by M', then define x to be the support vertex adjacent to w in G' or the other end of the edge in M' incident with w. Note that the degree of x is at least two. If w is a leaf in G' and x has degree two, then x is not a support vertex in G since G is not a path. Define $M = M' \cup \{wx\}$ if w is a leaf in G' and $\deg_G(x) = 2$, and define $M = M'$ for otherwise. So M is a matching that saturates no support vertices of G.

Let F defined on $V(G)$ obey M. Define S_u to be a subset of $F(v)$ with size $|F(v)| - 1$ such that $|S_u \cap F(x)| \leq 1$ (we assume that $F(x)$ is the empty set when x is not defined). Define $F' : V(G') \to 2^{[14]}$ by letting $F'(w)$ be a subset of $S_u - F(x)$ with size $|S_u| - 1$, and $F'(z) = F(z)$ for every vertex z in G' other than w. So F' obeys M', and there is an F'-avoiding pair (f'_1, f'_2) of G' with respect to I'. It is easy to define an F-avoiding pair (f_1, f_2) of G with respect to I' such that $f_i(w)$ is a subset of $f'_i(w) - S_u$ with size $|f'_i(w)| - 1$ and $f_i(u)$ contains S_u for $i = 1, 2$. ■

Given nonnegative integers α, β, an (α, β)-*star* is the graph obtained from $K_{1,\alpha+\beta}$ by subdividing β edges. A *spider* is a tree that has exactly one vertex, denoted by the *central vertex*, of degree at least three.

Lemma 4.7 *If G is a subcubic spider with the central vertex v, then G is $(M, \{v\})$-tractable for some matching M.*

Proof. By Lemma 4.6, it suffices to show that every (α, β)-star with central vertex v, where $\alpha + \beta = 3$, is $(\emptyset, \{v\})$-tractable. Let F obey $\emptyset$. Let $V(G) = \{v, x_i, y_j, y'_j : 1 \leq i \leq \alpha, 1 \leq j \leq \beta\}$ and $E(G) = \{vx_i, vy_j, y_jy'_j : 1 \leq i \leq \alpha, 1 \leq j \leq \beta\}$. We may assume that $|F(x)| = 6 - 2\deg_G(x)$ for every vertex x in G. If $(\alpha, \beta) = (3, 0)$, then let S_v be a subset of $[14]$ of size three such that S_v intersects $F(x_i)$ for $1 \leq i \leq 3$. If $(\alpha, \beta) = (2, 1)$, then let S_{y_1} be a subset of $F(y_1)$ with size three, S_v be a subset of $[14] - S_{y_1}$ of size two such that S_v intersects $F(x_1)$ and $F(x_2)$. If $(\alpha, \beta) = (1, 2)$, then let S_v be a subset of $F(x_1)$ with size one, and S_{y_j} a subset of $F(y_j) - S_v$ with size 3 for $j = 1, 2$. If $(\alpha, \beta) = (0, 3)$, then let S_{y_j} be a subset of $F(y_j)$ with size 3 for $j = 1, 2, 3$. Then it is easy to define an F-avoiding coloring f such that for every vertex x, $f(x)$ contains S_x (if S_x is defined) and $f(x)$ has size $8 - \deg_G(x) - 1_{\{v\}}(x)$ by first assigning $f(v)$ and then assigning $f(x_i)$ and $f(y_j)$ for $1 \leq i \leq \alpha$ and $1 \leq j \leq \beta$, and then assigning $f(y'_j)$ for $1 \leq j \leq \beta$. Hence, (f, f) is an F-avoiding pair of G with respect to $\{v\}$. ■

Lemma 4.8 *Let G be a connected subcubic graph which is not a path or a spider. Assume that G has a cut-edge v_2w such that $G - v_2w$*

contains a path $P = v_1v_2...v_{|P|}$ as a component with $|P| = 3$ or 4. Let $G' = G - P$. If G' is (M', I')-tractable for some matching M' of G' and independent set I' of G', then G is $(M, I' \cup \{v_2\})$-tractable for some matching M of G.

Proof. When w is a leaf in G' or w is saturated by M', we denote x by the support vertex adjacent to w in G' or the other end of the edge in M' incident with w. Since G is not a path or a spider, and M' does not saturate any support vertex in G', we know that when x has degree two and w is a leaf in G', x is not a leaf and not a support vertex in G. Define $M = M' \cup \{wx\}$ if w is a leaf in G' and x has degree two, otherwise, define $M = M'$. So M is a matching that does not saturate any support vertex in G.

Let F defined on G obey M, and let Y be the set of leaves adjacent to w. In the remaining of the proof, $F(x)$ is the empty set when x is not defined. If $|P| = 4$, then define S_{v_2} to be a subset of $F(v_1) - F(x)$ with size $|F(v_1)| - 3$, and S_{v_3} to be a subset of $F(v_4) - S_{v_2}$ with size $|F(v_4)| - 1$; if $|P| = 3$, then define S_{v_2} to be a smallest subset of $F(v_1) \cup F(v_3)$ such that $|S_{v_2} \cap F(v_1)| \geq |F(v_1)| - 3$, $|S_{v_2} \cap F(v_3)| \geq |F(v_3)| - 3$ and $|(S_{v_2} \cup F(w)) \cap (\bigcup_{y \in Y} F(y) \cup F(x))| \leq 1$. Note that S_{v_2} exists since $|\bigcup_{y \in Y} F(y) \cup F(x)| \leq 7$.

Define F' on $V(G')$ by assigning $F'(z) = F(z)$ for $z \neq w$, and assinging $F'(w)$ a subset of $(S_{v_2} \cup F(w)) - (\bigcup_{y \in Y} F(y) \cup F(x))$ with size $|S_{v_2} \cup F(w)| - 1$. So F' obeys M', and hence there is an F'-avoiding pair (f'_1, f'_2) of G' with respect to I'. And then it is easy to obtain an F-avoiding pair (f_1, f_2) of G with respect to $I' \cup \{v_2\}$ by modifying f'_1 and f'_2. Note that w has degree two in G', so $w \notin I'$ and $I' \cup \{v_2\}$ is an independent set in G. ■

Lemma 4.9 *Let t be a nonnegative integer, and let G be the graph obtained from a path $u_1u_2u_3u_4v_1v_2...v_{2t}u_5u_6u_7u_8$ by attaching one leaf u'_i on each vertex u_i for $3 \leq i \leq 6$. Then G is $(\emptyset, \{u_3, u_6\})$-tractable.*

Proof. Let F obey the empty set, so $F(u_1)$ is disjoint from $F(u_2)$ and $F(u_7)$ is disjoint from $F(u_8)$. Define S_{u_4} to be a subset of $F(u'_4)$ with size $|F(u_4)| - 2$, and for each $1 \leq i \leq t$, define $S_{v_{2i}}$ to be a subset of $(S_{v_{2i-2}} \cup F(v_{2i-1})) - F(v_{2i})$ with size $|S_{v_{2i-2}} \cup F(v_{2i-1})| - 2$, where v_0 is u_4. Furthermore, define S_{u_3} to be a subset of $F(u'_3) - S_{u_4}$ with size $|F(u'_3)| - 3$, S_{u_2} to be a subset of $F(u_1) - S_{u_3}$ with size $|F(u_1)| - 1$, S_{u_5} to be a subset of $F(u'_5) - S_{v_{2t}}$ with size $|F(u'_5)| - 2$, S_{u_6} to be a subset of $F(u'_6) - S_{u_5}$ with size $|F(u'_6)| - 3$, and S_{u_7} to be a subset of $F(u_8) - S_{u_6}$ with size $|F(v_8)| - 1$. Then it is easy to find an F-avoiding coloring f of G by first assigning $f(u_5)$ a superset of S_{u_5} with size 5, and then assigning $f(v_j)$ a subset of $[14] - (F(v_j) \cup f(v_{j+1}) \cup S_{j-1} \cup S_{j+1})$

containing S_j (let $S_k = \emptyset$ for undefined S_k, where $j-1 \leq k \leq j+1$) with size $8 - \deg_G(z) - 1_{\{u_3,u_6\}}(z)$ from $j = 2t$ down to -1, where $v_{-1} = u_3$ and $v_{2t+1} = u_5$, and then assigning $f(z)$ by a superset of S_z (if S_z is defined) for other vertices z with size $8 - \deg_G(z) - 1_{\{u_3,u_6\}}(z)$. Hence, (f, f) is an F-avoiding pair with respect to $\{u_3, u_6\}$. $\blacksquare$

Lemma 4.10 *Let G be a subcubic graph such that no support vertex has degree two. Let $v \in V(G)$ and $N_G(v) = \{v', u, w\}$, where v' is a leaf and u, w are of degree two. If $G' = G - \{v, v'\}$ is (M', I')-tractable for some matching M' and independent set I', then G is $(M' \cup \{uu', ww'\}, I' \cup \{v\})$-tractable, where u' and w' is the neighbor of u and w other than v, respectively.*

Proof. Let $M = M' \cup \{zz' : (z, z') = (u, u') \text{ or } (w, w'), \deg_G(z') = 2\}$. Note that u and w are leaves in G', so M' does not saturate u' and w'. And since no support vertex has degree two in G, M is a matching of G saturating no support vertices in G.

Let F defined on $V(G)$ obey M. Define S_v to be a subset of $[14]$ with size 4 such that S_v intersects $F(u)$, $F(w)$ and $F(v')$ with $|(S_v \cup F(u)) \cap F(u')| \leq 1$ and $|(S_v \cup F(w)) \cap F(w')| \leq 1$. Define F' on $V(G')$ by assigning $F'(u)$ a subset of $(F(u) \cup S_v) - F(u')$ with size $|F(u) \cup S_v| - 1$, and assigning $F'(w)$ a subset of $(F(w) \cup S_v) - F(w')$ with size $|F(w) \cup S_v| - 1$, and define $F'(z) = F(z)$ for every other vertex z in G'. Hence F' obeys M', and there is an F'-avoiding pair (f'_1, f'_2) of G' with respect to I', and then it is easy to obtain an F-avoiding pair (f_1, f_2) of G with respect to $I' \cup \{v\}$ such that $f_i(v) = S_v$ by modifying f_i, where $i = 1, 2$. $\blacksquare$

Recall that the family $\mathcal{H}$ of graphs consists of $H_{a,b}$ for all even integers $a, b \geq 4$.

Lemma 4.11 *Let G be a good graph such that no component of G is a path or a graph in $\mathcal{H}$. Then G is (M, I)-tractable for some matching M and independent set I.*

Proof. We shall prove this lemma by doing induction on $|V(G)|$. The base case follows from Lemma 4.7. Assume the lemma holds for every good proper subgraph of G containing no path or a graph in $\mathcal{H}$ as a component, but G is not (M, I)-tractable for any M and I. So G is connected and G is not a spider. The following claim is an immediate consequence of Lemmas 4.5 and 4.6.

Claim 1: If v is a support vertex of degree 2 adjacent to a nonsupport vertex, then $G - \{v, v'\}$ is a graph in $\mathcal{H}$, where v' is the leaf adjacent to v.

Claim 2: No support vertex of degree 2 is adjacent to a support vertex of degree 3.

Proof Claim 2: Suppose that there is a support vetex of degree two adjacent to a support vertex of degree three. Then there is a cut-edge e in G such that one component C of $G-e$ is a path of order 4. By Lemmas 4.5, 4.1 and 4.8, G is (M, I)-tractable for some M and I unless $G-C$ is a path of even order or a graph in $\mathcal{H}$. If $G-C$ is a path of even order, then G is either a spider or in $\mathcal{H}$ by Claim 1, a contradiction. And if $G-C$ is a graph in $\mathcal{H}$, then by Claim 1, either there is another cut-edge e' such that one component C' of $G-e'$ is a path of order four but $G-C'$ is not a graph in $\mathcal{H}$, or G is the graph mentioned in Lemma 4.9. So no support vertex of degree two adjacent to a support vertex of degree three. $\square$

Claim 3: No support vertex is of degree 2.

Proof of Claim 3: Let v be a support vertex of degree 2, and let $N_G(v)=\{v', w\}$, where v' is a leaf. Since C is a not path, w is not a support vertex by Claim 2. Hence, every support vertex of degree 2 is adjacent to a leaf and a nonsupport vertex. Together with Claim 1, there is no support vertex of degree 2. $\square$

Similarly, by taking advantage of Lemmas 4.5 and 4.8, we have the following Claim.

Claim 4: Every support vertex is adjacent to exactly one leaf.

In addition, the following claim follows from the bipartiteness of G, Claim 3 and Lemma 4.10.

Claim 5: No support vertex is adjacent to two vertices of degree two. Consequently, every support vertex is of degree three and adjacent to exactly one leaf and at least one vertex of degree three.

Let X be the subgraph of G induced by vertices of degree three. By (G1), X has maximum degree at most two, so X is a disjoint union of paths and cycles.

Claim 6: X is a disjoint union of paths.

Proof of Claim 6: Suppose that some component of X is a cycle. Since G is connected, G is obtained from an even cycle by attaching leaves on every vertex of it by (G2) and (G3). We shall obtain a contradiction by showing that G is $(\emptyset, \emptyset)$-tractable. Write $V(G)=\{v_i, u_i : 0 \leq i \leq 2k-1\}$ and $E(G)=\{v_iv_{i+1}, v_iu_i : 0 \leq i \leq 2k-1\}$, where the index is computed under modulo $2k$, for some positive integer k. Let F obey the empty set. Give an orientation on the edges of the cycle $v_0v_1...v_{2k-1}v_0$ such that every vertex in the cycle has in-degree and out-degree one. By Lemma 4.4, for every $0 \leq i \leq 2k-1$, we can pick a subset S_i of $F(u_i)$ with size $|F(u_i)|-2$ such that S_i is disjoint from $S_{i-1} \cup S_{i+1}$. Again by Lemma 4.4, for every $1 \leq i \leq 2k$, we can pick a subset T_i of $[14]-\bigcup_{j=i-1}^{i+1} S_j$ with size $5-|S_i|$. Then it is easy to define an F-avoiding coloring f of G such that $|F(z)| = 8 - \deg_G(z)$ for every vertex z in G and $f(v_i) = S_i \cup T_i$ for every $0 \leq i \leq 2k-1$. So (f, f) is an F-avoiding pair of G, and hence G is $(\emptyset, \emptyset)$-tractable. $\square$

The following claim is an immediate consequence of (G2), (G3), (G4) and Claim 5.

Claim 7: If P be a component in X, then the following hold.

1. If $|V(P)| \geq 3$, then every vertex in P is a support vertex in G.
2. If $|V(P)| = 2$, then P contains a support vertex in G.
3. If $|V(P)| = 1$, then the vertex in P is a nonsupport vertex of degree 3 in G.

Let $X_1, X_2, ...$ be the components of X. If X_i is a path of order two, then let $x_i \in X_i$ be a support vertex in G. Let I be the subset of $V(X)$ consisting of x_i, for each component X_i with $|V(X_i)| = 2$, and the ends of every other component of X. So I is an independent set on G, and every vertex in I has degree three in G. We shall show that G is $(\emptyset, I)$-tractable. Let G' be the graph obtained from G by deleting all leaves, so every support vertex in G has degree two in G'. Note that G' has minimum degree at least two by Claim 5.

Claim 8: There exists an orientation O_1 of G' such that every vertex has in-degree and out-degree at least one, and if $v \in V(G')$ with $\deg_{G'}(v) = 3$ is a nonsupport vertex in G, and v adjacent to a support vertex in G, then v is a head of an edge whose tail is a nonsupport vertex in G, and v is a tail of an edge whose head is a nonsupport vertex in G.

Proof of Claim 8: Let Y be the graph obtained from G' by deleting every edge that is incident with one support vertex in G and one nonsupport vertex in G of degree three. By Lemma 2.6, there is an orientation O of Y such that every vertex v has in-degree and out-degree at least $\lfloor \deg_Y(v)/2 \rfloor$. Let $J = \{u \in V(G') : u$ is a support vertex in G, $N_Y(y)$ contains a nonsupport vertex of degree three$\}$ is an independent set in G by (G4), so J is an independent set. Note that if $u \in J$, then u is incident with an edge in $E(G') - E(Y)$ by Claim 7. Define an orientation O_1 of G' by assigning the direction of each edge in $E(G') \cap E(Y)$ the same direction as in O, and assigning each edge in $E(G') - E(Y)$ a direction such that every vertex in J has the in-degree and the out-degree one. If $u \in V(G')$ is a support vertex in G, then either $\deg_Y(u) = 2$ or $u \in J$, so u has in-degree and out-degree one in O_1. If $u \in V(G')$ is a nonsupport vertex in G, then $\deg_Y(u) \geq 2$ by (G3), so u has in-degree and out-degree at least one in O_1. Furthermore, if $v \in V(G')$ is a nonsupport vertex with $\deg_{G'}(v) = 3$, and v is adjacent to a support vertex in G, then v is an end of some X_i with $|V(X_i)| = 2$, so the other two neighbors of v are of degree 2 in G. In other words, $\deg_Y(v) = 2$, and the two neighbors of v in Y are nonsupport vertices by Claim 5. Therefore, v is a head of an edge whose tail is a nonsupport vertex in G, and v is a tail of an edge whose head is a nonsupport vertex in G. $\square$

Let O_2 be the orientation of G' obtained from O_1 by reversing the direction of each edge. Define D_1 and D_2 to be the digraph whose underlining graph is G' equipped with orientation O_1 and O_2, respectively.

Let F defined on $V(G)$ obey the empty set, and we may assume that $|F(v)| = 6 - 2\deg_G(v)$. Since X is a union of paths, we can write $V(G) = \{v_i : 1 \leq i \leq |V(G)|\}$ such that every vertex in X has at most one neighbor with smaller index, and the index of every vertex in X is smaller than the index of any vertex not in X. Note that the ordering of $V(G)$ is also an ordering of $V(G')$. Now, for $i = 1$ and 2, we consecutively define $r_i : V(D_i) \to \mathbb{N}$ and T_i : {support vertices of $G\} \to 2^{[14]}$ according to the above ordering of $V(G')$ and the following rules:

1. Assume that v is the vertex in some component X_k of X with $|V(X_k)| = 1$. If $\deg^+_{D_i}(v) = 1$, then define $r_i(v) = 6$; otherwise, define $r_i(v) = 2$.
2. Assume that $|V(X)| \geq 2$, and v is an end of a component of X, and v is in I. So v is a support vertex in G, and we let v' be the leaf adjacent to v. Define $r_i(v) = 3$. If v is pointed by a vertex of degree 3, then define $T_i(v) = \emptyset$; otherwise, v is pointed by a vertex of degree 2, and we define $T_i(v)$ to be a subset of $F(v') - T_i(u)$ with size 2, where u is the neighbor of v such that $T_i(u)$ is defined (if any).
3. Assume that $|V(X)| \geq 2$, and v is an end of a compoenent of X, but v is not in I. So v is an end of a component X_k of X with $|V(X_k)| = 2$. Assume that v is not a support vertex in G. If v is pointed by a vertex of degree 3, then define $r_i(v) = 6$; otherwise, define $r_i(v) = 4$.
4. Assume that $|V(X)| \geq 2$, and v is an end of a compoenent of X, but v is not in I. Assume that v is a support vertex in G, and let v' be the leaf adjacent to v. If v is pointed by a vertex of degree 3, then define $r_i(v) = 3$ and $T_i(v) = \emptyset$; otherwise, define $r_i(v) = 2$, and $T_i(v)$ to be a superset of $F(v')$ with size 5 such that $T_i(v) \cap F(w) \neq \emptyset$, where u is the neighbor of v of degree 2.
5. Assume that $|V(X)| \geq 2$, and v is an internal vertex of a component of X. So v is a support vertex in G, and we let v' be the leaf adjacent to v. Define $r_i(v) = 3$, and $T_i(v)$ to be a subset of

$F(v') - T_i(u)$ with size 2, where u is the neighbor of v such that $T_i(u)$ is defined (if any).

6. Define $r_i(v) = 6$ for every vertex v with $\deg_G(v) = 2$.

It is clear that $r_1(v) + r_2(v) + |T_1(v)| + |T_2(v)| = 16 - 2\deg_G(v) - 2 \cdot 1_I(v)$ for every vertex v in G' (i.e. every non-leaf in G). Define $L_i : V(D_i) \to 2^{[14]}$ by letting $L_i(v) = [14] - (F(v) \cup \bigcup_{u \in N_{G'}[v]} T_i(u))$ for every vertex v in G'. So $|L_i(v)| \geq r_i(v) + \sum_{u \in N^+_{D_i}} r_i(u)$ for every vertex v in G'. By Lemma 4.4, there are F-avoiding colorings g_1 and g_2 of G' such that $g_i(v) \subseteq L_i(v)$ and $|g_i(v)| \geq r_i(v)$ for $i = 1, 2$, so $|g_1(v)| + |g_2(v)| \geq r_1(v) + r_2(v)$. Define $f_i(v) = g_i(v) \cup T_i(v)$ for $i = 1, 2$ and $v \in V(G')$ (we let $T_i(v)$ be the empty set if $T_i(v)$ is not defined), so $|f_1(v)| + |f_2(v)| \geq r_1(v) + r_2(v) + |T_1(v)| + |T_2(v)| = 16 - 2\deg_G(v) - 2 \cdot 1_I(v)$ for $v \in V(G')$. In addition, for every support vertex v adjacent to the leaf v', $|f_i(v) - F(v')| \leq r_i(v) + 1$, and the equality holds only when $r_i(v) = 2$ and $|T_i(v)| = 5$. So $|f_i(v) - F_i(v')| \leq 3$ for every support vertex v and the leaf v' adjacent to it, and hence we can define $f_i(v')$ to be a subset of $[14] - (f_i(v) \cup F(v'))$ such that $|f_i(v')| = 7$. As a result, (f_1, f_2) is an F-avoiding pair of G with respect to I. This completes the proof. ■

5 Penetrations and cooperations

In this section, we will prove Lemmas 2.2, 2.3 and 2.4, and it completes the proof of lemmas in section 2.

Proof of Lemma 2.2: Let $C = v_1v_2...v_k$, where k is odd. If $k = 1$, then let $B = (N_G(C), \emptyset)$. If $k \geq 5$, then let $B = (N_G(C), \{u_1u_2, u_{k-1}u_k : u_i \in N_G(v_i) \cap N_G(C)$ for $i = 1, 2, k-1, k\})$. Then $(B, \emptyset)$ penetrates C by Lemma 4.1. Note that B is loopless since G is triangle-free.

When $k = 3$, let B_1 and B_2 be graphs such that $V(B_1) = V(B_2) = N_G(C)$ and $E(B_1) = \{x_1x_2 : x_i \in N_G(v_i) \cap N_G(C), i = 1, 2\}$, $E(B_2) = \{x_2x_3 : x_i \in N_G(v_i) \cap N_G(C), i = 2, 3\}$. Then it is easy to check that $(B_1, \emptyset)$ cooperates with $(B_2, \emptyset)$. □.

Proof of Lemma 2.3: First, we shall prove this lemma when C is a path of even order. Let $C = v_1v_2...v_{2k}$, and B_1, B_2 be graphs with $V(B_1) = V(B_2) = N_G(C)$. Now we define $E(B_1)$ and $E(B_2)$. If $\deg_G(v_i) \leq 2$ for some $1 \leq i \leq 2k$ and $i \neq 2, 2k-1$, then define $E(B_1) = E(B_2) = \{u_ju_{j+1} : u_j \in N_G(v_j) \cap N_G(C), j = 1, 2k-1\}$. Note that $\deg_{B_i}(v) \leq 2n_C(v)$ for every vertex v in B_i. By statement 12 of Lemma 4.2, both $(B_1, \emptyset)$ and $(B_2, \emptyset)$ penetrate C. So we may assume that $\deg_G(v_i) = 3$ for $i \neq 2, 2k-1$.

Let $N_G(v_1)-C=\{u_1,u_1'\}$, $N_G(v_{2k})-C=\{u_{2k},u_{2k}'\}$ and $N_G(v_i)-C=\{u_i\}$ for $2\le i\le 2k-1$. Note that u_2 and u_{2k-1} may not exist, and we just ignore it when we mention it in the remaining of this proof if it does not exist. If $k=1$, then define $E(B_1)=E(B_2)=\{xy: x\in N_G(v_1)\cap N_G(C), y\in N_G(v_2)\cap N_G(C)\}$. If $k=2$ and $N_G(v_1)\cap N_G(v_4)=\emptyset$, then define $E(B_1)=\{u_1u_2,u_1'u_2,u_3u_4,u_3u_4',u_1u_4,u_1u_4'\}$ and $E(B_2)=\{u_1u_2,u_1'u_2,u_3u_4,u_3u_4',u_1u_4,u_1'u_4\}$. If $k=2$ and $N_G(v_1)\cap N_G(v_4)\ne\emptyset$, then $|N_G(v_1)\cap N_G(v_4)|\le 1$ since there is no f-rainbow L_0, and we assume that $u_1'=u_4'$ and define $E(B_1)=E(B_2)=\{u_1u_2,u_1'u_2,u_3u_4,u_3u_4',u_1u_4\}$. Then $(B_1,\emptyset)$ and $(B_2,\emptyset)$ are desired by statements 1 and 2 of Lemma 4.2.

Now we assume that $k\ge 3$. Let $s=-\infty$ if $N_G(v_1)\cap N_G(v_4)=\emptyset$; otherwise, let $s=\max\{i\in\mathbb{Z}: N_G(v_{2j+1})\cap N_G(v_{2j+4})\ne\emptyset, 0\le j\le i\}$. Note that either $s=-\infty$ or $0\le s\le k-2$. If $s\ge 0$, then let $u_1\in N_G(v_4)$. If $s=k-2$, then let $u_{2k}\in N_G(v_{2k-3})$.

Case 1: If $s\ge 1$ and $n_C(u_1')\ge 2$, then define $E(B_1)=E(B_2)=\{u_1u_2,u_1'u_2,u_{2k-1}u_{2k},u_{2k-1}u_{2k}'\}$, and define J by the following rule:

- $J=\{u_1,u_1',u_3\}$ if $n_C(u_1)=n_C(u_1')=n_C(u_3)=2$;
- $J=\{u_1,u_3\}$ if $n_C(u_1)=n_C(u_1')=n_C(u_3)=3$ and $u_1'\in N_G(v_{2k-1})$;
- J is any independent subset of $\{u_1,u_1',u_3\}$ of size 2 containing a vertex of n_C-value three, otherwise.

Then $(B_1,\{J\})$ and $(B_2,\{J\})$ are desired unless $u_1'\in N_G(v_{2k-1})$ and $n_C(u_1')=2$, by statements 7 and 8 of Lemma 4.2. Note that a similar augument works for the case that $s\ge 0$, $n_C(u_1')\ge 2$, $u_1'\notin N_G(v_{2k-1})$ and $n_C(u_3)\ge 2$ unless $u_3\in N_G(v_{2k-1})\cup N_G(v_{2k})$ and $n_C(u_3)=2$. Similarly, the lemma holds if $s\ge 0$, $n_C(u_1')=3$, $u_1'\in N_G(v_{2k-1})$, and $n_C(u_3)\ge 2$. So we may assume that either $s\le 0$, or $s\ge 1$ and one of the following: $n_C(u_1')\le 1$ or $n_C(u_1')=2$ with $u_1'\in N_G(v_{2k-1})$.

Case 2: If $s=k-2$, then define $E(B_1)=E(B_2)=\{u_1u_2,u_1'u_2, u_{2k-1}u_{2k},u_{2k-1}u_{2k}',u_1'u_{2k}',u_{2j-1}u_{2j}: 2\le j\le k-1\}$. Note that $s=k-2\ge 1$, so either $n_C(u_1')\le 1$ or $n_C(u_1')=2$ with $u_1'\in N_G(v_{2k-1})$, and hence $u_1'\ne u_{2k}'$. Then $(B_1,\emptyset)$ and $(B_2,\emptyset)$ are desired by statement 3 of Lemma 4.2.

Case 3: If $s=k-3$, then define $E(B_1)=E(B_2)=\{u_1u_2,u_1'u_2, u_{2j-1}u_{2j},u_{2k-1}u_{2k},u_{2k-1}u_{2k}',u_{2k-3}u_{2k},u_{2k-3}u_{2k}': 2\le j\le k-2\}$. Then $(B_1,\emptyset)$ and $(B_2,\emptyset)$ are desired by statement 5 of Lemma 4.2.

Case 4: If $0\le s\le k-4$, and if $u_1'\notin N_G(v_{2s+6})$ when $s=0$, then $k\ge 4$ and $u_1'\notin N_G(v_{2s+6})$ since either $n_C(u_1')\le 1$ or $n_C(u_1')=2$ with $u_1'\in N_G(v_{2k-1})$ when $s\ge 1$, and then we define $E(B_1)=\{u_1u_2,u_1'u_2,u_{2j-1}u_{2j},u_{2k-1}u_{2k},u_{2k-1}u_{2k}',u_1'u_{2s+6},u_{2s+3}u_{2s+6},u_{2s+5}u_{2s+6}: 2\le j\le s+1\}$, and define $E(B_2)$ later. Then $(B_1,\emptyset)$ penetrates C by statement 6 of Lemma 4.2, and at most one vertex v in B_1 has degree more than $2n_C(v)$. Moreover, if such v exists, then v is adjacent to a vertex u in B_1 with $\deg_{B_1}(u) \leq 2n_C(u) - 1$, and v is adjacent to a vertex v_i in C with even index $i = 2s+6 \geq 4$; if $\deg_{B_1}(v) \geq 2n_C(v)+1$ and $n_C(v) \geq 2$ then $n_C(v) = 2$ and v is adjacent to v_{2k-1}.

Case 5: If $s = -\infty$, then define $E(B_1) = \{u_1u_2, u'_1u_2, u_{2k-1}u_{2k}, u_{2k-1}u'_{2k}, u_1u_4, u'_1u_4, u_3u_4\}$, and define $E(B_2)$ later. Similarly, $(B_1, \emptyset)$ penetrates C by statement 4 of Lemma 4.2, and at most one vertex v in B_1 has degree more than $2n_C(v)$, and once such v exists, v is adjacent to a vertex with the index 4, and $n_C(v) \geq 2$ implies $n_C(v) = 2$ and v is adjacent to v_{2k-1}.

If there is no vertex v such that $\deg_{B_1}(v) = 2n_C(v) + 1$, or the vertex v with $\deg_{B_1}(v) = 2n_C(v)+1$ satisfies $n_C(v) = 3$ in Case 4 and Case 5, then we further define $E(B_2) = E(B_1)$.

Case 6: Assume that $s = 0$ and $u'_1 \in N_G(v_6)$, and if $n_C(u_3) \geq 2$, then $n_C(u_3) = 2$ and $u_3 \in N_G(v_{2k-1}) \cup N_G(v_{2k})$. Note that it is the remaining case that $E(B_1)$ has not been defined. We consider the following subcases.

- When $k \geq 5$ (so $u_3 \neq u_8$) and $u_5 \neq u_8$, define $E(B_1) = \{u_1u_2, u'_1u_2, u_3u_8, u_5u_8, u_7u_8, u_{2k-1}u_{2k}, u_{2k-1}u'_{2k}\}$, then $(B_1, \emptyset)$ are desired by statement 10 of Lemma 4.2. Note that every vertex v other than u_8 in B_1 has degree at most $2n_C(v)$. And $\deg_{B_1}(u_8) \geq 2n_C(u_8)+1$ implies that $n_C(u_8) \neq 2$ or $n_C(u_8) = 2$ with $u_8 \in N_G(v_2) \cup N_G(v_{2k-1})$. We further define $E(B_2) = E(B_1)$ if $\deg_{B_1}(u_8) \leq 2n_C(u_8)$ or $n_C(u_8) = 3$.

- When $k \geq 4$ and $u_5 = u_8$, let $t = \max\{i \in \mathbb{Z} : F(v_{2j+5}) \subseteq F(v_{2j+8}), 0 \leq i \leq j\}$, so $0 \leq t \leq k-4$. If $t = 2k-4$, then we assume that $u_{2k} \in N_G(v_{2k-3})$. Define $E(B_1) = \{u_1u_2, u'_1u_2, u_5u_6, u_{2i+7}u_{2i+8}, u_{2k-1}u_{2k}, u_{2k-1}u'_{2k} : 0 \leq i \leq t-1\} \cup T$, where T is defined as follows:

  - $T = \{u_3u'_{2k}\}$ if $t = k-4$;
  - $T = \{u_{2k-3}u_{2k}, u_{2k-3}u'_{2k}\}$ if $t = k-5$;
  - $T = \{u_3u_{2t+10}, u_{2t+7}u_{2t+10}, u_{2t+9}u_{2t+10}\}$ if $0 \leq t \leq k-6$.

  Notice that B_1 does not have loops since $n_C(u_3) \geq 2$ implies that $n_C(u_3) = 2$ and $u_3 \in N_G(v_{2k-1}) \cup N_G(v_{2k})$, and hence $(B_1, \emptyset)$ penetrates C by statement 11 of Lemma 4.2. Also, every vertex v other than u_{2t+10} in B_1 has degree at most $2n_C(v)$, and $\deg_{B_1}(u_{2t+10}) \geq 2n_C(u_{2t+10}) + 1$ only if $0 \leq t \leq k-6$. And either $n_C(u_{2t+10}) \neq 2$, or $n_C(u_{2t+10}) = 2$ with $u_{2t+10} \in N_G(v_2) \cup N_G(v_{2k-1})$. We further define $E(B_2) = E(B_1)$ if $\deg_{B_1}(u_{2t+10}) \leq 2n_C(u_{2t+10})$ or $n_C(u_{2t+10}) = 3$.

- When $k = 4$ and $u_5 \notin N_G(v_8)$, define $E(B_1) = E(B_2) = \{u_1u_2, u_1'u_2, u_5u_8, u_5u_8', u_7u_8, u_7u_8'\}$, then $(B_1, \emptyset)$ and $(B_2, \emptyset)$ are desired by statement 9 of Lemma 4.2.
- When $k = 3$, define $E(B_1) = E(B_2) = \{u_1u_2, u_1'u_2, u_3u_6, u_3u_6', u_5u_6, u_5u_6'\}$, then $(B_1, \emptyset)$ and $(B_2, \emptyset)$ are desired by statement 5 of Lemma 4.2.

Consequently, $E(B_1)$ are defined in all cases.

In fact, if B_2 is not defined, then $k \geq 3$ and there is an even number $p \notin \{2, 2k\}$ and a vertex $w_1 \in N(v_p)$ in B_1 such that the following hold:

- $\deg_{B_1}(w_1) = 2n_C(w_1) + 1$;
- $n_C(w_1) \leq 2$;
- if $n_C(w_1) = 2$, then $w_1 \in N_G(v_p) \cap N_G(v_{2k-1})$ or $w_1 \in N_G(v_p) \cap N_G(v_2)$;
- if $v \in V(B_1) - \{w_1\}$, then $\deg_{B_1}(v) \leq 2n_C(v)$;
- if $v \in V(B_1) - \{w_1\}$ and $\deg_{B_1}(v) = 2n_C(v)$, then $N_G(v) \cap V(C) \subseteq \{u_1, u_2, u_{2k-1}, u_p\}$.

Let $x_i = v_{2k+1-i}$ for $1 \leq i \leq 2k$, so $x_1x_2...x_{2k}$ is the same path as $v_1v_2...v_{2k}$ but the order of the index of vertices are reversed. Now, for those case that $E(B_2)$ have not been defined, we define other boundry-graphs B_1', B_2' for the path $x_1x_2...x_{2k}$ with $V(B_1') = V(B_1) = V(B_2') = V(B_2)$ by applying the same rules as we defined B_1, B_2. The same argument shows that either $E(B_1')$ and $E(B_2')$ can be defined, or $E(B_1')$ can be defined in a way such that every vertex v except one vertex $w_2 \in N(v_{2k+1-q})$ with some even number $q \notin \{2, 2k\}$ in B_1' has degree at most $2n_C(v)$ such that the following hold:

- $\deg_{B_2}(w_2) = 2n_C(w_2) + 1$;
- $n_C(w_2) \leq 2$;
- if $n_C(w_2) = 2$, then $w_2 \in N_G(v_{2k+1-q}) \cap N_G(v_2)$ or $w_2 \in N_G(v_{2k+1-q}) \cap N_G(v_{2k-1})$;
- if $v \in V(B_1') - \{w_2\}$ and $\deg_{B_1}(v) = 2n_C(v)$, then $N_G(v) \cap V(C) \subseteq \{u_2, u_{2k-1}, u_{2k}, u_{2k+1-q}\}$.

Hence, we are done if both $E(B_1)$ and $E(B_2)$ or both $E(B_1')$ and $E(B_2')$ can be defined, so it remains to consider the case that $E(B_2)$ and $E(B_2')$ are not defined. We claim that $(B_1, \emptyset)$ and $(B_1', \emptyset)$ are desired. Notice that it is enough to prove that $w_1 \neq w_2$. Since $n_C(w_1) \leq 2$ and $n_C(w_2) \leq 2$, we know that $w_1 = w_2$ implies that $w_1 \in N_G(v_2) \cap N_G(v_{2k-1})$, and it is a contradiction as $p \neq 2$. This proves the lemma when C is a path of even order.

When C is a graph in $\mathcal{H}$. Let $V(C) = \{x_i, y_j : 1 \leq i \leq a, 1 \leq j \leq b\}$ and $E(C) = \{x_i x_{i+1}, y_j y_{j+1}, x_2 y_2 : 1 \leq i \leq a-1, 1 \leq j \leq b-1\}$ for some even positive integers a and b. Let $V(B_1''') = V(B_2''') = N_G(\{x_i : 1 \leq i \leq a, i \neq 2\}) \cap N_G(C)$. Define $E(B_1''')$, $E(B_2''')$, $\mathcal{J}_1'''$, and $\mathcal{J}_2'''$ as in the case of path of even order. Note that since $N_G(x_2) \cap N_G(C) \cap (V(B_1''') \cup V(B_2''')) = \emptyset$, we have that for $i = 1, 2$, if there exists $x \in J_i'''$ for some $J_i''' \in \mathcal{J}_i'''$ with $n_C(x) = 3$, then $|\bigcup_{v \in J_i'''} N_{B_i'''}(v)| \leq 1$. And then define B_1 and B_2 be graphs on $N_G(C)$ such that $E(B_i) = E(B_i') \cup \{wz : w \in N_G(y_{b-1}) \cap N_G(C), z \in N_G(y_b) \cap N_G(C)\}$. For each $i = 1, 2$ and each $J_i''' \in \mathcal{J}_i'''$, if $J_i''' \cap (N_G(y_{b-1}) \cup N_G(y_b)) = \emptyset$, then define $J_i = J_i'''$; otherwise, define J_i to be an independent subset of J_i''' of size 2 containing a vertex with n_C-value 3 such that $|\bigcup_{v \in J_i} N_{B_i}(v)| \leq 3$. Then $(B_1, \mathcal{J}_1)$ and $(B_2, \mathcal{J}_2)$ penetrates C by Lemma 4.3, where $\mathcal{J}_1$ and $\mathcal{J}_2$ is the collection of those J_1's and J_2's that we just defined, respectively. This completes the proof. $\square$

Proof of Lemma 2.4: Since f is a good 3-coloring, C is a good graph. By Lemma 4.11, there is a matching M and an independent set $I \subseteq \{v : \deg_C(v) = 3\}$ such that for every $F_1, F_2 : V(C) \to 2^{[14]}$ that obey M, there are F_i-avoiding colorings f_i of C, for $i = 1, 2$, such that $|f_1(v)| + |f_2(v)| = 16 - 2\deg_C(v) - 2 \cdot 1_I(v)$ for all $v \in V(C)$. Let $S = \{v'u' : v' \in N_G(v) \cap N_G(C), u' \in N_G(u) \cap N_G(C), \deg_C(v) = 1, u \in N_C(v)\}$. Let $T = \{x'y' : xy \in M, x' \in N_G(x) \cap N_G(C), y' \in N_G(y) \cap N_G(C)\}$. Define $B_1 = B_2 = (N_G(C), S \cup T)$. By Lemma 4.11, $(B_1, \emptyset)$ cooperates with $(B_2, \emptyset)$. $\square$

6 Concluding remarks

In this paper, we proved that $\chi_f(G) \leq 43/15$ for every triangle-free subcubic graph G. In fact, we proved the stronger statement that for every fractionally t-critical triangle-free subcubic graph G with $t \geq 8/3$, there is an independent set I and a function $f : V(G) \to 2^{[168]}$ such that $|f(v)| = 72 - 4\deg_G(v) - 4 \cdot 1_I(v)$ for every vertex v. Hence, $\chi_f(G) \leq 14/5$ if we can get rid of the error term $1_I(v)$ from the above equation. Recall that every vertex in I is adjacent to three vertices which receive the same color from a good 3-coloring. So one possible way to remove the error term is to prove that every fractionally 14/5-critical graph has a linear 3-coloring without any rainbow L_0. A linear coloring is a proper coloring such that each two color classes induce a graph of maximum degree at most 2. It is not hard to show that such a 4-coloring exists by using Liu and Yu's result about linear colorings [10]. On the other hand, Esperet et al. [2] showed that determining whether a subcubic graph admits a linear 3-coloring is NP-complete in general, so the condition for fractionally critical is crucial.

In contrast with linear coloring, we conjecture that every subcubic graph other than K_4 has a proper 3-coloring such that each two color classes induce a disjoint union of cacti, where a cactus is a graph whose blocks are edges and cycles. At some time when we tried to simplify the proof, we thought that the positive answer of the conjecture might be helpful.

Finally, we note that Heckman and Thomas [8] also conjectured that every planar triangle-free subcubic graph has fractional chromatic number at most 8/3. In section 2, we investigate the structure of fractionally t-critical graphs with $t \geq 8/3$, so it might be helpful in dealing with this conjecture.

Acknowledgement. The author thanks Professor Robin Thomas for many suggestions for preparing this paper.

References

[1] W. Abbas, M. Egerstedt, C.-H. Liu, R. Thomas and P. Whalen, *$(5,2)$-configuration on minimum degree at least two $K_{1,6}$-free graphs*, manuscript, 2011.

[2] L. Esperet, M. Montassier and A. Raspaud, *Linear choosability of graphs*, Discrete Math. 308 (2008), 3938–3950.

[3] S. Fajtlowicz, *On the size of independent sets in graphs,* Cong. Numerantium 21 (1978), 269–274.

[4] D. Ferguson, T. Kaiser and D. Král', *The fractional chromatic number of triangle-free subcubic graphs,* arXiv:1203.1308v2.

[5] S. Gutner and M. Tarsi, *Some results on (a,b)-choosability,* Discrete Math. 309 (2009) 2260–2270.

[6] H. Hatami and X. Zhu, *The fractional chromatic number of graphs of maximum degree at most three,* SIAM J. Discrete Math. 23 (2009) 1762–1775.

[7] C. C. Heckman and R. Thomas, *A new proof of the independence ratio of triangle-free cubic graphs,* Discrete Math. 233 (2001) 233–237.

[8] C. C. Heckman and R. Thomas, *Independent sets in triangle-free cubic planar graphs,* J. Combin. Theory Ser. B 96 (2006) 253–275.

[9] F. Kardoš, D. Král' and J. Volec, *Fractional colorings of cubic graphs with large girth,* SIAM J. Discrete Math. 25 (2011) 1454–1476.

[10] C.-H. Liu and G. Yu, *Linear colorings of subcubic graphs,* arXiv:1206.5348v1.

[11] L. Lu and X. Peng, *The fractional chromatic number of triangle-free graphs with* $\Delta \leq 3$, Discrete Math. 312 (2012), 3502–3516.

[12] M. Richardson, *Solutions of irreflexive relations*, Ann. of Math. 58 (1953) 573–580.

[13] E. R. Scheinerman and D. H. Ullman, Fractional Graph Theory. A Rational Approach to the Theory of Graphs, Wiley-Intersci. Ser. Discrete Math. Optim., John Wiley & Sons, Inc., New York, 1997.

[14] W. Staton, *Some Ramsey-type numbers and the independence ratio,* Trans. Amer. Math. Soc. 256 (1979), 353–370.